


REPORT

COOLCITY Index 2026

Adaptive potential of 11,041 European Union cities



Editorial team

The report presents analyses based on data from the COOLCITY Index 2026 database, covering 11,041 cities in the European Union.

The COOLCITY Index 2026 Report was created as part of the project "Use of remote sensing for management of blue-green infrastructure in the process of city adaptation to climate change" (LIFECOOLCITY) spanning 2023-2029, by a consortium: **MGGP Aero (leader)**, **City of Wrocław**, **University of Warsaw**, **University of Łódź**, **Ingenieurgesellschaft Prof. Dr. Sieker mbH**.

The project is co-financed by the European Union under the LIFE programme and the National Fund for Environmental Protection and Water Management.

Author: MGGP Aero © (2026) MGGP Aero Ltd.

Graphic design: Monika Krzesiak

Layout: Jarosław Aleksander

How to cite

MGGP Aero. (2026). **COOLCITY Index 2026 Report: Adaptive potential of 11,041 European Union cities**. coolcity.eu

Database

Kopeć, D., Takmadżan, A., Jedliński, J., Żmuda, D., Tarkowska, A., Migas, R., Krok, M., Łochowski, M., & Sławik, Ł. (2026). The COOLCITY Index 2026 for 11,041 cities in the European Union [Data set]. Zenodo. <https://doi.org/10.5281/zenodo.20638922>

COOLCITY Index 2026 data are available under the CC BY-NC-SA license.

AUTHORS:



DSc
Dominik Kopeć



Łukasz
Sławik



Zuzanna
Kunert



Aleksandra
Krzywicka



dr Agata
Tarkowska



Robert
Migas



Alicja
Śniadach



Aram
Takmadżan



Martyna
Wietecha



Jan
Jedliński



Dominik
Żmuda



Jakub
Charyton



Wiktor
Zwierzyński



Monika
Krzesiak



Jarosław
Aleksander

Table of contents

About the report	4
------------------	---

Foreword	5
----------	---

Project context	8
------------------------	---

COOLCITY as a response to the needs of cities	11
---	----

Adaptative challenges	12
-----------------------	----

What COOLCITY Index can offer cities	13
--------------------------------------	----

Tools for cities	14
------------------	----

COOLCITY Index in the European Union	15
---	----

Key findings	16
--------------	----

One Index, many realities	17
---------------------------	----

Four potential regions	18
------------------------	----

Spectrum of city experiences	20
------------------------------	----

Unequal starting points on the path to adaptation	21
---	----

Performance Index	22
-------------------	----

Blue-green city profiles	23
---------------------------------	----

• Northern Europe	25
-------------------	----

• Western Europe	29
------------------	----

• Central-Eastern Europe	33
--------------------------	----

• Southern Europe	37
-------------------	----

Regional leaders	41
-------------------------	----

COOLCITY regional leaders	41
---------------------------	----

Regional leaders Performance Index	42
------------------------------------	----

Best practices from the blue-green backyard	43
---	----

Shared challenges and adaptation opportunities	46
--	----

COOLCITY Index in EU countries	47
---------------------------------------	----



Wrocław - the first COOLCITY in Europe	100
---	-----

Wrocław in the COOLCITY Index 2026	101
------------------------------------	-----

Nature-based adaptation	103
-------------------------	-----

Data-driven management	104
------------------------	-----

COOLCITY Index Methodology	106
-----------------------------------	-----

A city in COOLCITY	106
--------------------	-----

How is the COOLCITY Index (CCI) created?	107
--	-----

Indicators	108
------------	-----

Data sources and bibliography	110
-------------------------------	-----



About the report

Climate change is an enormous challenge for European cities. In the face of increasing frequency of flash floods, climbing exposure to droughts and heatwaves, and ongoing ecosystem degradation, local governments must know how to build resilience and ensure a good quality of life for their residents.

Our ambition is to provide key knowledge and universal tools to support cities in adapting to climate change based on nature. That is why we created this report and the [COOLCITY Index platform](#), presenting our analyses of the state of blue-green infrastructure in over 11,000 cities across the European Union.

What does it present?

The COOLCITY Index 2026 report is an in-depth analysis of the environmental state of European cities in 2025. The report characterises distinctive city profiles in various EU regions and examines the situation in each of the 27 EU countries. It diagnoses the challenges cities face today, reveals their causes, and suggests how to address them. The report and platform will be updated and published annually, enabling monitoring of changes over time.

Who is it for?

The report is aimed at anyone wishing to deepen their knowledge and make decisions based on reliable and objective data. This includes decision-makers at European, regional, and national levels; scientists, educators, media representatives, NGOs, and anyone interested in the impact of climate change on cities and their inhabitants.

How was it created?


The report interprets data available on the [COOLCITY Index 2026 platform](#). It is a comprehensive database on urban areas of the EU, containing administrative, demographic, geographic, and climatic information from publicly available official sources such as satellite data from ESA, NASA, Google DW, EUROSTAT, IMERG, IPCC, and GloH2O, as well as our proprietary COOLCITY indicators measuring cities' adaptive potential based on their environmental resources.

Foreword

Today's increasing interaction of urbanization processes and climate change within the material structure of cities - especially the high proportion of sealed surfaces, loss of biodiversity, and the density of population and infrastructure concentration - amplifies our exposure to extreme events. Under such conditions, episodes of high temperatures, periods of water scarcity, or intense rainfall directly impact the functioning of urban systems and public health. At the same time, it is precisely in cities that management processes occur, enabling the integration of spatial, environmental, and health policies to strengthen resilience ^[1].

Current knowledge indicates that the configuration of the urban environment is directly linked to residents' health risks. High surface temperatures, intensified by the urban heat island effect and the predominance of impermeable, heat-retaining materials, increase health burdens, especially in densely built-up areas. Simultaneously, appropriately selected features of the city's spatial layout - including the share of greenery, water availability, building form, and mobility organisation - allow for influence on exposure levels to these threats ^[2].

Nature-based solutions and blue-green infrastructure should therefore be regarded as key elements of climate resilience strategies. Systems combining vegetation, water, and ecological functions can simultaneously increase retention, improve water quality, reduce heating, and support urban habitat functioning ^[3]. This approach aligns with the conclusion of the Pontifical Academy of Sciences workshop "Resilience of People and Ecosystems under Climate Stress." Nature-based solutions at the building and city scale were presented as contributing to reduced energy consumption, supporting natural cooling, humidification, and air purification, and strengthening the long-term resilience of ecosystems ^[4].






The significance of this approach goes beyond a single study, as the workshops resulted in a set of recommendations forming the basis of the Vatican's memorandum for the 2022 UN summit in Sharm el-Sheikh, with a clear shift in emphasis from mitigation alone to adaptation and resilience building.

Thus, investments in urban nature should be understood as multifunctional interventions, important both for climate adaptation and for public health and the sustainability of urban systems. In this context, comparing cities solely through the lens of individual sectoral indicators is methodologically insufficient. Analyses conducted for hundreds of European cities show that population density, building morphology, and access to natural areas coexist with variations in urban heat island intensity, pollution, and health indicators. It is particularly important that compact cities can combine lower per capita emissions with limited access to green spaces and deterioration of certain local environmental parameters if densification is not accompanied by appropriate quality and scale of green infrastructure ^[5].

The significance of the COOLCITY Index 2026 platform lies precisely in the need to synthesise this complexity into a comparable, scalable, and operationally useful form. The value of this approach lies not only in aggregating data for over 11,000 urban areas but above all in establishing a common diagnostic model for assessing the environmental adaptive potential of cities. In conditions of growing demand for evidence-based decisions, synthetic tools provide an important, organised, and complementary addition to local diagnosis.





Effective adaptation requires the intentional multifunctionality of green infrastructure, understood as designing solutions that simultaneously benefit water balance, thermal conditions, biodiversity, and residents' well-being. The adaptive potential of green spaces is spatially and qualitatively uneven. Global analyses of cooling generated by urban greenery show that thermal benefits depend on both the quantity and quality of greenery, with their distribution varying significantly between cities and world regions^[6]. Meanwhile, epidemiological modelling indicates that increased greening can lower ambient temperatures, leading to a statistically significant reduction in heat-related mortality^[7]. Consequently, assessing adaptive potential should encompass not only the area of green spaces but also their condition, structure, interconnections, and ecological functionality^[8].

Incorporating biodiversity into urban resilience analyses is crucial, as it supports ecosystem services, enhances adaptive capacity, and delivers measurable benefits for human health and well-being, while simultaneously facing spatial competition pressures, especially where management integration is insufficient^[9]. This means that diagnostics of blue-green infrastructure should be conducted systemically – as an analysis of the relationships between water, vegetation, temperature, permeability, habitat continuity, and urbanisation pressure.

In Europe, this approach is particularly necessary due to the strong variation in climatic conditions and urban development trajectories. Water deficits, overheating, densification pressures, and topographical constraints occur with varying intensity across different parts of the continent, so interpreting adaptive potential requires simultaneously considering comparative metrics and regional context. The COOLCITY Index 2026 report addresses this need by combining a common assessment method with the ability to interpret results through the lens of diverse environmental profiles.

A particular value of the COOLCITY project is that it does not stop at the measurement stage. The integration of the Index, View, and Design tools aligns with the increasingly prevalent logic in urban research: from integrated diagnosis, through priority identification, to the selection of interventions tailored to local constraints and resources. This approach enhances the practical utility of environmental data in planning and management, where climate adaptation must take the form of spatial, investment, and operational decisions.

In this sense, the COOLCITY Index 2026 is not just a compilation of results but a tool that organises knowledge about the environmental conditions of urban resilience. It demonstrates that adaptive capacity stems from measurable environmental properties and city structure. At the same time, it reminds us that a climate-resilient city is also a healthier city, more efficiently managed, and better integrated with natural processes.

Therefore, the report should be read not only as a diagnosis of the current state but also as a comparative and analytical instrument supporting the design of future actions. Where indicators show high environmental resources, it can strengthen relationships and knowledge exchange between cities and the identification of good practices. Where it reveals deficits, it can form the basis for more precise intervention planning. Nature-based adaptation is linked to a model of city shaping in which climate safety, public health, and ecological integrity are treated as interdependent goals of urban policy.



Barbara Widera, PhD, DSc, Arch.

Associate Professor, Faculty of Architecture, Wrocław University of Science and Technology,
Member of the EU Mission Board on Climate Change Adaptation

Project context

Vulnerable Cities of the Future

The future of Europe is urban. By 2021, the European Union had approximately 447 million inhabitants, of whom around 337 million lived in urban and suburban areas. This means that urban spaces are the daily environment for three-quarters of the EU population [10]. It is estimated that this number will rise to 83% by 2050 [11].

The quality of urban space therefore determines the quality of life for Europeans. A city is not just a layout of buildings, roads, and technical infrastructure. It is a human living environment. Its form decides whether residents have access to shade, water, greenery, places to rest, and daily contact with nature. It affects physical and mental health, ease of movement, opportunities for recovery, social activity, and a sense of security. The WHO emphasises that interventions related to urban greenery can support people's health and wellbeing, including mental health, prevention of cardiovascular diseases, and physical activity [12].

75%

mieszkańców UE w 2021
to mieszkańcy miast



83%

mieszkańców UE w 2050
to będą mieszkańcy miast




Project context

Increasing Climate Threats

The importance of the urban environment and its impact on human well-being is growing, especially under increasing climate pressure. Recent years have been among the warmest on record [13], and Europe is the fastest-warming continent – with temperature increase rates twice the global average. And heatwaves claim the deadliest toll among climate threats [14].

In Europe, an average of over 175,000 people die annually due to extreme temperatures. The WHO indicates that with ongoing climate change, urbanisation, and an ageing population, these numbers may only rise. Historical records confirm this trend: of the 23 most severe heatwaves to hit Europe since 1950, 19 have occurred since 2000 [15].

The effects of climate change are felt particularly strongly in cities. Population concentration, a high proportion of impermeable surfaces, and dense development exacerbate challenges related to heatwaves, local overheating, drought, rapid stormwater runoff, and flooding [16]. Therefore, appropriate quality of the natural environment becomes one of the key conditions for ensuring the safety of city residents today.



TEMPERATURE ANOMALIES WORLDWIDE AND IN EUROPE


Source: C3S/ECMWF • <https://climate.copernicus.eu/esotc/2025/why-europe-warming-so-quickly>

Adaptation Based on Nature

The relationship between humans and the urban environment amid increasing climate threats is becoming a prerequisite for equitable access to good living conditions. Deficits in greenery and biodiversity, surface impermeability, and adverse water and thermal conditions are interconnected problems that affect the resilience of cities and the health of their inhabitants, who must adapt to living in a warmer, more extreme climate.

Therefore, blue-green infrastructure (BGI) should be regarded as strategic urban infrastructure. Climate adaptation in the European Union is increasingly focused on implementing nature-based solutions (NBS). The EU Adaptation Strategy [17] promotes the use of BGI in cities as an effective tool to mitigate the impacts of heatwaves, floods, and droughts, and emphasises the importance of NBS as multifunctional actions that can provide greater social and environmental benefits than traditional technical solutions [18].

Adaptation requires the conscious development of the city's natural resources. BGI works in two ways: it increases the city's resilience to climate change while simultaneously improving residents' living conditions through the ecosystem services it provides. It is a valuable asset that also requires protection. Without proper planning, maintenance, and preservation of space for nature, it can degrade, causing the city to lose part of its natural capacity to mitigate climate change effects. Adaptation thus requires balance: humans benefit from nature's services but must also create conditions in which urban nature can thrive.



COOLCITY as a response to the needs of cities

COOLCITY Index 2026

City managers, planners, and residents need current, clear, and comparable knowledge about the urban environment. Without it, it is difficult to identify where the city has the greatest adaptive potential, where deficits accumulate, and which areas require further diagnosis. Thus, the key question is not only how to improve quality of life in the city but also how to obtain data and translate it into decisions.

The answer to this need is the COOLCITY Index 2026, a platform and report presenting data on the adaptive potential of European cities. At its core is the **COOLCITY Index (CCI)** – a synthetic measure describing the state of blue-green infrastructure. The index is calculated for over 11,000 urban areas, with values on a scale from 0 to 10 (the higher the value, the better), allowing for a consistent analysis and placement on a common map of Europe.




IMAGE OF EUROPEAN CITIES SEEN FROM SATELLITE:
PIWNICZNA ZDRÓJ, STOCKHOLM, ATHENS
Source: modified Copernicus Sentinel data from 2025

Many challenges, one measure


The CCI index is created based on satellite data and GIS. These allow us to analyse “cities” as urban areas defined by a single, consistent method. CCI enables systematic monitoring of many cities on the same scale, according to the same principles and based on comparable data. It does not replace local diagnosis but allows for a quick recognition of the overall state of green infrastructure and adaptive challenges.

CCI integrates five key elements of the urban environment: **surface permeability**, **vegetation state**, **biodiversity**, **water conditions**, and **thermal conditions**. These components are in continuous interaction, mutually enhancing or limiting the city's resilience. Together, they create a synthetic overview of whether the urban environment supports nature-based adaptation.



Adaptive challenges

COOLCITY Index



Surface permeability

0-10

This is the degree of surface sealing in the city and its ability to infiltrate rainwater. High index values indicate a large proportion of biologically active surfaces that promote water retention and the functioning of urban ecosystems, while low values indicate a high degree of imperviousness and a limited capacity for natural water circulation.



Vegetation state

0-10

Indicates vegetation coverage and its condition, with particular emphasis on tall vegetation. High index values indicate well-developed and lush vegetation that supports the city's resilience to climate change and the preservation of ecosystem services. Low values indicate a deficit or deteriorated state of greenery requiring protective or restorative actions.



Biodiversity

0-10

Describes the degree of diversity and naturalness of vegetation, as well as the connectivity of ecological corridors in the city. High indicator values show the presence of diverse forms of vegetation that support biodiversity conservation, while low values indicate a lack of networked natural structures and limited conditions for the functioning of living organisms.



Water conditions

0-10

They define the city's ability to retain water and maintain water-related ecosystems. They take into account the land's retention potential and the proportion of wetland areas.

High values of the indicator show favourable water conditions and greater resilience of the city to the effects of drought and heavy rainfall, while low values indicate limited retention.



Thermal conditions

0-10

This refers to the city's susceptibility to overheating and the severity of the urban heat island, taking into account surface temperature and the radiative properties of the ground. High index values indicate favourable thermal conditions and greater resilience of the city to heatwave effects, while low values suggest an increased risk of overheating.


What the COOLCITY Index offers European cities

The European challenge framework

The greatest value of the COOLCITY Index Report and platform lies in the ability to view cities as a unified European system. The COOLCITY Index addresses questions that cannot be seen from the perspective of a single city: where environmental conditions are relatively best, where deficits accumulate, how small, medium, and large cities differ, and how a city's conditions compare to those of its country, region, or to the entire Europe. This allows the platform not only to present data on the urban environment but also to organise the spatial arrangement of adaptive challenges, translating complex environmental data into clear information about the adaptive potential of cities.

Global perspective, local action

The CCI indicator enables an overview of the whole spectrum of perspectives, from a global view of the state of Europe's urban ecosystems to detailed local-level analyses, providing information on the internal diversity of BGI resources within any city. COOLCITY Index, together with COOLCITY View and COOLCITY Design products, forms three mutually complementary tools that support local governments at every stage of the adaptation process – from environmental diagnosis, through strategic planning, to the implementation of specific nature-based actions.



Tools for cities

COOLCITY consists of three tools based on the CCI: the freely accessible [COOLCITY Index](#) and the commercial solutions [COOLCITY View](#) and [COOLCITY Design](#). They support cities in diagnosing environmental conditions, planning climate change adaptation, and designing nature-based solutions.

The [COOLCITY Index](#) allows for an initial, comprehensive diagnosis of the adaptive potential of European cities and identifies the main challenges related to blue-green infrastructure.


More detailed environmental diagnosis and planning support are provided by the View and Design tools:

[COOLCITY View](#) is a strategic tool based on satellite data, comprising a spatial data platform and an expert report. It supports diagnosis and monitoring of environmental conditions, identification of priority areas requiring action, and climate adaptation planning. For analysed spatial units, it also indicates the recommended direction of blue-green infrastructure actions: protection, redevelopment, or introduction of new solutions.

[COOLCITY Design](#) is a decision support system based on detailed aerial data, consisting of a spatial data platform and an expert report. It assists cities in making decisions regarding the implementation of NBS. It enables analysis of various implementation scenarios, considering ecosystem services, costs, and environmental benefits resulting from planned interventions. This helps assess which actions can bring the greatest adaptive and environmental effect in specific locations.

You can learn more about COOLCITY on the [platform coolcity.eu](https://platform.coolcity.eu).

**COOLCITY
Index**



Satellite data

**COOLCITY
View**



Satellite data

**COOLCITY
Design**



Aerial data

COOLCITY Index in the European Union

The COOLCITY Index, covering 11,041 European cities, is one of the largest data sources enabling the assessment of the environmental adaptive capacity of urban areas in the European Union. The inclusion of such a large number of cities allowed for comparative analyses, pattern identification, and the formulation of conclusions regarding the state of the blue-green infrastructure (BGI) in 2025 and the challenges facing European cities.

COOLCITIES IN NUMBERS



11,041
cities




289 million
inhabitants



134,000 km²
Urban area

COOLCITY Index



Key findings

The analysis shows that European cities vary significantly in terms of BGI status and adaptation needs, both within individual countries and geographic regions, as well as between groups of different population sizes. Dividing cities into regional profiles helps compare centres with similar conditions and highlights places with the greatest deficits, even in regions with relatively favourable BGI conditions in Europe.

[Read more about profiles](#) →

The highest adaptive potential is found in very small cities, under 10,000 inhabitants, regardless of region and profile.

[Read more](#) →

90.5 million (31.3%)
of Europe's population lives in urban areas with an **unfavourable environmental profile**.

49.5 million (17.1%)
people live in resilient cities with a **favourable profile** and very good BGI condition.

[Read more](#) →

In over 800 EU cities, the environmental condition is very good and there is a high potential for climate change adaptation (CCI >7). The highest proportion of such cities is in Finland (almost 60%).

More than 1,000 European cities have very low adaptive capacity based on BGI (CCI <4).

The greatest challenge is in Cyprus, where all urban areas have very low CCI values (maximum 3.4 CCI).

[Read more](#) →

The strongest correlation with a city's adaptive potential measured by the CCI index is population density and geographic location.

[Read more](#) →

One index, many realities

What is the current adaptive potential of European countries?


We examined the COOLCITY Index values in cities across EU countries to assess how prepared they are for current and future climate challenges.

The distribution of the index across Europe reveals significant variation in cities' adaptive potential. A clear spatial pattern emerges: as you move northwards in Europe, CCI values gradually increase.

The highest adaptive potential was recorded in Finland, with an average index value of 7.1. Alongside Latvia (CCI 7.0), these are the only countries with a very high adaptive potential. At the opposite end are Cyprus, with an average of 2.1, and Malta, with 3.8. Both countries have low potential, indicating greater challenges in adaptation using NBS resources.

These results highlight a general trend observed continent-wide: cooler climates and conditions typical of northern Europe more often support building resilience based on nature. They illustrate the different realities in which individual countries operate.

COOLCITY Index



Four regions of potential


Regions in the European Union

Cities face unequal starting points on the path to adaptation due to their location on the map of Europe, so it is crucial to view them through the lens of regional conditions.

Dividing European cities into regions allows for comparing their adaptive potential under similar geographical conditions, while also capturing differences in green infrastructure resources within the regions themselves.

The report highlights 4 regions:

- Northern Europe
- Western Europe
- Central-Eastern Europe
- Southern Europe




How varied is the state of the BGI across EU regions?

The most favourable pattern of adaptive potential is visible in Northern Europe. The region includes 567 cities, with an average CCI value of 6.6, the highest in the entire EU. As many as 96% of northern cities achieve a score equal to or above the European average of 5.5 CCI, and 26% reach values above 7 CCI. The highest proportion of cities with such high adaptive potential is found in Finland (almost 60%). The low variability of results in northern countries shows that cities in the region have uniformly high potential linked to favourable environmental conditions.

A different picture is seen in Southern Europe. The average indicator value for cities in the region (3771) is 4.9. Only 1 in 3 southern cities reaches the EU average score. At the same time, this region shows the greatest contrasts – alongside areas with relatively high potential, there are cities with clearly weaker environmental resources. Cities in the region range from 0.7 up to 8.4 CCI. An unfavourable BGI state is reached by 20% of cities here (CCI below 4). The most internally diverse countries are Greece (from 0.7 to 8.4) and Italy (from 0.8 to 7.9), due to the great geographical diversity within the country (from coastal cities to those located in higher mountain areas).

Western and Central-Eastern Europe form an intermediate pattern. They include 4952 and 1751 cities respectively, with an average indicator value of 5.8 in both regions. Cities in these parts of Europe generally score above average but do not show as clearly a favourable picture as in the North. In the West, 70% of cities achieve a CCI value equal to or above the EU average, and in Central-Eastern Europe 67%. The most diverse country is Poland, where the CCI range spans from 4.4 to 8.5 (including the highest individual city score in the entire EU).

CCI VARIATION IN EU COUNTRIES



COOLCITY Index

0-1 1-2 2-3 3-4 4-5 5-6 6-7 7-8 8-9 9-10

Low potential

High potential

25% (first quartile)

average

individual cities

75% (third quartile)

Spectrum of city experiences

How does city size shape the potential in European countries?


Let's take a closer look at how adaptive potential depends on city population size and the share of built-up area in its spatial structure.

The division of cities by population classes clearly shows that larger cities generally have lower potential than smaller centres. The most favourable are the smallest cities, below 10,000 inhabitants, with an average CCI value of 5.8. The score gradually decreases: to 5.5 in cities of 10-20 k population, down to 4.9 in cities with over 100k residents. The problem of low urban environmental resilience is therefore most pronounced where the population is highest.

	VERY SMALL < 10k inhabitants	SMALL 10 - 20k inhabitants	MEDIUM 20 - 100k inhabitants	LARGE 100 - 500k inhabitants	VERY LARGE > 500k inhabitants
CCI	5.8	5.5	5.2	4.9	4.9
built-up area coverage	27.0%	30.1%	34.0%	37.2%	37.4%

This pattern is most clearly visible in France, where potential gradually falls from 5.7 CCI in very small cities, through 4.5 in medium-sized cities, to 3.5 in metropolises over 500,000 inhabitants. Meanwhile, neighbouring Germany shows less variability in BGI regardless of city size (respectively: 6.3, 6.0, 5.7).

In the largest cities, urbanisation pressures accumulate: strong densification, higher share of built-up areas, and less availability of BGI resources. The average built-up coverage rises from 27.0% in very small towns to 37.4% in very large ones (over 500,000 inhabitants). The scale of challenges increases moving southwards. Southern cities are clearly more compact and intensively built-up (34.8% built-up area), while northern cities have 12% less built-up area (22.8%). Urbanisation pressure is particularly visible in Cyprus and Spain, exacerbating adaptation difficulties. Large Southern cities face significantly greater challenges than small Northern cities.



VARIAION OF CCI IN EU COUNTRIES BY CITY SIZE CLASSES

Unequal starting points on the path to adaptation

What shapes the condition of the urban environment?

We already know that the adaptive potential of European urban centres depends on geographic location and population size. However, this does not provide the full picture. Cities do not start from the same baseline, and their blue-green infrastructure resources used to build climate resilience are not equal. What else influences the adaptive potential measured by the COOLCITY Index?

The key variables are:

population density – the higher the density, the lower the index values, indicating a greater share of built-up areas and pressure on blue-green infrastructure;

distance from the equator (latitude) – the further north, the better the conditions for nature-based adaptation, linked to favourable climatic conditions, i.e. lower temperatures and higher precipitation;

and terrain diversity – which creates barriers to development and helps preserve space for vegetation.

Therefore, the adaptive potential of cities is influenced not only by planning decisions but also by geographical conditions over which local governments have limited control. Hence, simply comparing the entire population of European cities without considering these factors can lead to oversimplified conclusions.

A lower COOLCITY index value should not be interpreted as weaker city actions. It often indicates a more challenging starting point. This is why assessing adaptive potential should consider both the current state of the urban environment and the context in which the city operates.




CHART OF THE IMPACT OF CITY ATTRIBUTES ON MODELLED CCI VALUES (SHAP)

Performance Index

Who performs best within their conditions

The value of the **COOLCITY Index** alone is not sufficient for a full interpretation of a city's adaptive actions. Therefore, we created an additional indicator to help assess whether cities effectively use their resources and develop high-quality blue-green infrastructure.

The **Performance Index (PI)** shows the difference between the COOLCITY Index value for a given city and the average value of other cities with similar socio-geographical conditions. It reflects the benefits gained from applying BGI in cities adaptation relative to local conditions and opportunities.

The index models the impact of socio-geographical conditions on the city's adaptive potential and allows assessment of how the state of BGI in a given city differs from that of cities developing under similar geographical conditions and levels of urbanisation. This way, it is possible to identify not only which cities have high adaptive potential but also which use their resources better than expected based on their conditions. A positive **PI** value indicates a better-than-expected result, while a negative value may indicate untapped potential, yet to be tapped by the city's governance.

How varied are **PI** values across European regions? The greatest range of the index is seen in Southern Europe, in very small towns, where values range from -1.8, which negatively assesses the current state of BGI compared to similar cities in the region, up to 1.5, showing how these centres effectively use their resources and shape a relatively high adaptive potential. The smallest variation in the **PI index** occurs in the Northern region, where cities range from -1.1 to 1.0 relative to the expected adaptive potential.

The expected COOLCITY Index is influenced by:

climatic conditions

geographical location


terrain formation

population density

↑ PERFORMANCE INDEX

shows how much better or worse a city performs than expected, based on results from cities with similar conditions

shows the benefit of applying blue-green infrastructure in city adaptation



Interpretation of the Performance Index

< 0.0
the city performs worse than cities with similar conditions

0.0
the city's result aligns with cities of similar conditions

> 0.0
the city performs better than cities with similar conditions thanks to BGI

Blue-green city profiles

Blue-green city profiles allow for a more precise capture of the complexity of Europe's regional conditions, which do not conform to national borders. Similar challenge patterns may appear in cities across different parts of the region that share similar environmental conditions and urban characteristics.

Therefore, in each EU region, we have identified 3 groups of cities based on COOLCITY Index values and adaptive challenges, as well as population, population density, rainfall, temperature, and altitude.

Northern Europe

- BLUE-GREEN TOWNS OF THE NORTH
- RAINY LOWLAND CITIES
- URBAN BALTIC CENTRES

Western Europe

- SMALL SPONGE TOWNS
- NEUTRAL WESTERN CITIES
- WESTERN METROPOLISES UNDER PRESSURE


Central-Eastern Europe

- UPLAND CITIES OF GREEN ADVANTAGE
- LARGE CITIES WITH MODERATE CONDITIONS
- DENSE WARM CITIES

Southern Europe


- SOUTHERN PLACES OF RESPITE
- TYPICAL MEDITERRANEAN CITIES
- HEATED CENTRES OF THE SOUTH

City profiles enable a deeper understanding of inequalities in the state of the blue-green infrastructure (BGI) and the resulting adaptive potential of European cities. They organise the pattern of challenges in the region, where relatively **favourable** and **unfavourable** conditions for building resilience prevail. Knowing what shapes the potential of each city type allows us to effectively plan adaptive actions tailored to the specifics of each centre, rather than just the general characteristics of the entire region.




How to read city profiles?

Profile pages present a synthetic overview of the adaptive potential of cities in European regions.
To fully understand the information presented, it is helpful to follow four reading steps
- from general context to detailed results:



HOW TO READ THE DEVIATION CHART?

COOLCITY Index	6.6 (± 0.2)
Surface permeability	5.8 (± 0.5)
Vegetation state	6.5 (± 0.1)
Biodiversity	5.5 (± 1.0)
Water conditions	7.3 (± 0.5)
Thermal conditions	6.3 (± 0.2)



Direction and strength of deviation

The direction of the bar shows whether the result is higher or lower than the average. Its length is the z-score, meaning the longer the bar, the stronger the deviation from the average

Label and interpretation

The label indicates the value of the profile's mean difference from the region average, and the colour shows how to interpret it (positively or negatively)

Reference point


The zero axis represents the average indicator values in the region

Northern Europe

The Northern Europe region stands out in Europe for its very favourable conditions for the development of blue-green infrastructure. Lower air temperatures and relatively high rainfall create an environment where it is easier to maintain healthy vegetation and more stable water conditions.

There is also a less intensive urban development pattern here, with lower population density and a smaller share of sealed surfaces, which supports the preservation of biologically active and retention areas, resulting in high adaptive potential values. This means that BGI does not need to be rebuilt from scratch but can rely on existing resources. Almost one quarter of the region's population (4.4 million) live in cities characterised by very favourable BGI conditions.

The problem of urban overheating in Northern Europe is not as widespread as in warmer parts of the continent, so local thermal pressure spots are particularly visible here – mainly larger and more sealed urban centres. In these places, the region's cool climate is insufficient to effectively reduce the urban heat island effect, making additional BGI necessary. This city profile is home to 60% of the region's population.



18.5 million


urban
residents



567

cities

SHARE OF CITIES
IN BGI PROFILES
IN THE REGION:



Blue-green city profiles in the region

BLUE-GREEN TOWNS OF THE NORTH

● Example cities: Jūrmala, Oulu, Hillerød, Kiruna, Haapsalu, Druskininkai

- mostly the smallest population centres with low population density, where low urban pressure supports the preservation of natural resources
- CCI values for these cities are higher than the region average (6.6), showing that nature-based adaptation has particular potential here
- notable for high greenery per inhabitant and a large share of water and wetlands – the highest in the region and EU

Average indicators for profiles:

C 7.1

 15.2k inhabitants
1.1k people/km²

 **826 m²**
green space area per capita

 **5.6%**
share of **wetland ecosystems**

RAINY LOWLAND CITIES

● Example cities: Roskilde, Visby, Palanga, Valmiera, Rakvere, Mariehamn

- usually small in population but more densely populated and urbanised than “Blue-Green Towns of the North”
- more common in the southern part of the region, where higher population density is linked to increased sealed surface area
- this group's advantage is high rainfall, which helps maintain good conditions for greenery development and local ecosystems

Average indicators for profiles:

C 6.3

 15.1k inhabitants
1.5k people/km²

 **26.1%**
sealed surface area

 rainfall:
861 mm annually

URBAN CENTRES ON THE BALTIC SEA

● Example cities: Stockholm, Helsinki, København, Tallin, Rīga, Vilnius

- by far the largest cities in the region, highly urbanised, often located in coastal zones
- CCI value is on average 0.8 points lower than the region average (6.6), mainly due to greater sealing and lower vegetation cover
- despite the region's cool climate, these centres face local surface overheating problems, indicating their lower adaptive potential

Average indicators for profiles:

C 5.8

 135.9k inhabitants
2.6k people/km²


 **30.5%**
sealed surface area

 **surface temperature averaging 1.7°C**
higher than the region (32.1 °C)

Different conditions, a shared need for adaptation

What distinguishes each profile and how to enhance their potential?

COOLCITY INDICATORS



How to increase their adaptive potential?

Recommended direction for adaptive actions
based on nature and BGI for cities in this profile


**Protection of natural resources
and strengthening their continuity**
Increasing ecological connectivity
>19% improves the **biodiversity** index
by an average of **+0.5**

**Improvement of the quality and
diversity of blue-green infrastructure**
High vegetation cover >9% is
associated with an average **+0.6**
higher **green condition** index

**Implementation of nature-based
solutions**
Increasing the surface retention
capacity by 5% can improve water
conditions by **+0.6**

Residents of profiles in countries

Comparison of the number of residents of individual profiles in the region's countries



● BLUE-GREEN TOWNS OF THE NORTH

● RAINY LOWLAND CITIES


● URBAN CENTRES ON THE BALTIC COAST

Western Europe

Western Europe is the most urbanised region on the continent. It's nearly 5,000 cities are home to over 125 million people. In more than 600 centres, high building density and a large share of sealed surfaces limit the availability of biologically active areas, resulting in a low adaptive potential—one of the lowest in Europe. The problem of urban heat islands is particularly evident in these cities, intensifying adaptation challenges and creating adverse living conditions during heatwaves for 28% of residents.

The region's temperate climate, including relatively high rainfall, partially mitigates the effects of urbanisation, so the vast majority of people live in areas with average (57%) or good (14%) conditions for building resilience based on the BGI.

The most favourable conditions in the region are found in “Small sponge towns.” Thanks to a large share of permeable areas, their advantage is a high retention capacity. This allows them to more effectively retain rainwater, mitigate drought effects, and maintain more favourable thermal conditions than larger, more sealed cities.



125.7 million

urban
residents



4,952

cities

SHARE OF CITIES
IN BGI PROFILES
IN THE REGION:



Blue-green city profiles in the region

SMALL SPONGE TOWNS

● Example cities: Salzburg, Baden-Baden, Chamonix-Mont-Blanc, Spa, Killarney

- mostly smaller towns with low urban pressure, where a smaller share of built-up areas supports the preservation of natural resources
- good water retention capacity combined with high rainfall creates favourable conditions for vegetation growth in the region's cities
- these conditions translate into better greenery status and a higher adaptation potential compared to other cities in the region

Average indicators for profiles:

C 6.5



9.8k inhabitants
1.1k people/km²



62.0% surface retention capacity



Vegetation state +0.9
above the region average

NEUTRAL WESTERN CITIES

● Example cities: Hamburg, Bruges, Graz, Cork, Arnhem, Luxembourg

- cities of typical for the region size and population density, mainly located on lowland areas
- vegetation share is average here, but its condition is weaker than in other cities of the region
- achieve average resilience and BGI potential relative to the region and the entire EU, as indicated by moderate COOLCITY indicator values

Average indicators for profiles:

C 5.7



29.7k inhabitants
1.6k people/km²



27.9%
sealed surface area



71.4% of the area covered by vegetation,
including 64.9% with high vigour

WESTERN METROPOLISES UNDER PRESSURE

● Example cities: Paris, Brussels, Vienna, Dublin, Amsterdam, Antwerpen

- the largest and most densely populated cities in the region
- despite similar location and rainfall as “Neutral Western Cities”, they experience greater urban pressure
- combined with higher air temperatures, this results in unfavourable thermal conditions and CCI values below the regional and European average

Average indicators for profiles:

C 4.3



53.5k inhabitants
3.8k people/km²




44.9%
sealed surface area



surface temperature
3.1°C higher than the region (38.0 °C)

Different conditions, a shared need for adaptation

What distinguishes each profile and how to enhance their potential?



How to increase their adaptive potential?

Recommended direction for adaptive actions based on nature and BGI for cities in this profile



Protection of natural resources and retention potential

Improving retention capacity by 5% is associated with an average increase of +0.5 in the water conditions index



Improvement in the quality and diversity of blue-green infrastructure

In cities where ecological connectivity exceeds 86%, the biodiversity index increases on average by +0.8




Implementation of nature-based solutions

Limiting the share of high-temperature areas to 5% can improve thermal conditions by +0.7

Residents of profiles in countries

Comparison of the number of residents of individual profiles in the region's countries




Central-Eastern Europe

Central-Eastern Europe holds an intermediate position between the most favourable conditions for BGI development in Northern Europe and the more demanding conditions of the South.

The best conditions are primarily found in cities located in upland areas. Higher rainfall totals and greater retention capacity support maintaining good vegetation health and greater biodiversity, enabling these cities to achieve the highest adaptive potential values in the region. About 13% of the population lives in locations with favourable conditions. At the same time, more than half of the inhabitants live in large centres with moderate conditions. The relatively low share of sealed surfaces and high vegetation cover help maintain a balance between urban development and BGI functioning.

Higher sealing characterises southern cities in the region, so the problem of urban overheating is more pronounced there. It is precisely in these centres that BGI development can bring the greatest benefits, reducing the urban heat island effect and improving the resilience of the environment where nearly 14 million people live.



- POLAND
- CZECH REPUBLIC
- SLOVAKIA
- HUNGARY
- ROMANIA
- BULGARIA
- CROATIA
- SLOVENIA



51.5 million


urban
residents



1,751

cities

SHARE OF CITIES
IN BGI PROFILES
IN THE REGION:



- UPLAND CITIES OF GREEN ADVANTAGE
- LARGE CITIES WITH MODERATE CONDITIONS
- SEALED WARM CITIES

Blue-green city profiles in the region

UPLAND CITIES OF GREEN ADVANTAGE

- Example cities: Piwniczna-Zdrój, Karlovy Vary, Dubrovnik, Gabrovo, Celje
- generally the smallest and least densely populated cities, usually located in upland and mountainous areas with high rainfall totals
- in this group, CCI values are above the region average, showing that nature-based adaptation has particular potential here
- favourable environmental conditions, especially vegetation state and water conditions, visible in high retention capacity

Average indicators for profiles:

C 6.6

 10,000 inhabitants
1,200 people/km²

 **vegetation state +1.1**
above the region average

 rainfall: **914 mm** annually
63% surface retention capacity

LARGE CITIES OF MODERATE CONDITIONS

- Example cities: Warsaw, Prague, Bratislava, Zagreb, Ljubljana
- this group mainly includes the largest and most densely populated cities in the region, but with a relatively low share of impermeable surface per capita
- with moderate vegetation cover and typical regional green quality
- favourable thermal conditions in these cities mean urban pressure is felt less, and CCI values are only 0.3 lower than the region average

Average indicators for profiles:

C 5.7

 48,400 inhabitants
2,100 people/km²

 150.7 m² of sealed surface area per capita

 **70.6% of surface covered by vegetation,**
including 71.4% with high vigour

SEALED WARM CITIES

- Example cities: Sofia, Bucharest, Split, Szeged, Koper, Leszno
- mostly lowland, medium-sized cities, with average population density for the region
- high average annual air temperatures and significant surface sealing cause overheating problems in these centres
- adaptation using BGI is a greater challenge here compared to other cities in the region, reflected in the lowest COOLCITY indicator values

Average indicators for profiles:

C 4.8


 30,800 inhabitants
1,800 people/km²

 35.4%
sealed surface area

 **surface temperature**
4.5°C higher than the region

Different conditions, shared need for adaptation

What distinguishes each profile and how to enhance their potential?



How to increase their adaptive potential?

Recommended direction for adaptive actions based on nature and BGI for cities in this profile

Protection of natural resources and strengthening their continuity

An increase in vegetation richness by 0.2 corresponds to an average +0.6 higher biodiversity index

Improvement of the quality and diversity of blue-green infrastructure


High vegetation cover above 9% is associated with an average +0.6 higher green condition index

Implementation of nature-based solutions

Reducing the share of areas with high temperature to below 4% is linked to an average +0.7 higher thermal condition index

Residents of profiles in countries

Comparison of the number of residents of individual profiles in the region's countries



Southern Europe

Southern Europe is the most geographically diverse region in Europe, which is reflected in the greatest variability of BGI development conditions. Large variations in temperature, precipitation, and altitude mean that alongside cooler, wetter upland and mountainous towns, there are hot, highly urbanised Mediterranean cities where the development of high-quality BGI faces the greatest constraints and the potential for adaptation is the lowest on the continent.


The highest potential is retained by small towns located in upland and mountainous areas. Lower temperatures and higher rainfall support vegetation maintenance, enabling these towns to achieve the best adaptive indicators in the region. Such towns are home to 22% of the population. Nearly half of the region's inhabitants (46%) live in typical Mediterranean cities where urban pressure coexists with less favourable climatic conditions and achieve average BGI status. The greatest challenges are concentrated in the largest agglomerations. High sealing levels, very high population density, and hot, dry climate mutually reinforce the urban heat island effect. These cities have the lowest adaptive potential in the EU and include every third person in the region.

**93.4 million**urban
residents**3,771**

cities

SHARE OF CITIES
IN BGI PROFILES IN THE REGION:

30.2 million
438 cities



- SOUTHERN REFUGE TOWNS
- TYPICAL MEDITERRANEAN CITIES
- HEATED SOUTHERN CENTRES

-  PORTUGAL
-  SPAIN
-  ITALY
-  GREECE
-  CYPRUS
-  MALTA



Blue-green city profiles in the region

SOUTHERN RETREAT TOWNS

● Example cities: Sorrento, Santiago de Compostela, Sintra, Corfu

- towns with the lowest population and density in the region
- upland and mountainous centres with favourable climatic conditions providing lower temperatures and the highest rainfall totals among cities in the region and across the EU
- thanks to the highest CCI value in the region, cities of this profile stand out for their vegetation health and high water retention capacity

Average indicators for profiles:

C 5.8



11.2k inhabitants
1.4k people/km²



rainfall: 1,059 mm annually
57.5% surface retention capacity



64% share of high vigor vegetation

TYPICAL MEDITERRANEAN CITIES

● Example cities: Rome, Bilbao, Braga, Heraklion, Qormi

- mostly small but more densely populated towns with fairly high sealing
- usually located by the sea, lowland, with a higher average temperature (15.7°C) and lower rainfall totals than the region average
- although they have higher vegetation cover than the region average, their vegetation state and share of high biodiversity areas are low, resulting in lower climate resilience

Average indicators for profiles:

C 4.4



28k inhabitants
2.5k people/km²



38.4%
sealed surfaces



41% areas
with high biodiversity

HEATED SOUTHERN CENTRES

● Example cities: Barcelona, Lisbon, Athens, Nicosia, Valletta, Milan

- the largest cities in the region, heavily urbanised, with the highest population density in the EU and the highest sealing levels
- unfavourable thermal conditions and low rainfall combine here with low water retention capacity
- high urban pressure and intense surface overheating are linked to low adaptive potential of these centres – the CCI value is 2 points below the region average and the lowest in the entire EU

Average indicators for profiles:

C 2.9



68.9k inhabitants
4.8k people/km²




>55% sealed surfaces



20 % fewer low-temperature areas than the region average

Different conditions, shared need for adaptation

What distinguishes each profile and how to enhance their potential?



How to increase their adaptive potential?

Recommended direction for adaptive actions based on nature and BGI for cities in this profile



Protection of natural resources and strengthening their continuity

Increasing the share of areas with high biodiversity >18% corresponds to an average +0.5 higher biodiversity index



Improving the quality and diversity of blue-green infrastructure

Increasing high vegetation cover >7% is associated with an average +0.5 higher green condition index




Implementation of nature-based solutions

Increasing the share of areas with low temperature >4% corresponds to an average +1.0 higher thermal conditions index

Residents of profiles in countries

Comparison of the number of residents in each profile across the region's countries



● SOUTHERN PLACES OF RESPITE

● TYPICAL MEDITERRANEAN CITIES

● HEATED CENTRES OF THE SOUTH


COOLCITY regional leaders

How can we jointly build resilient cities?

Cities of the European Union form a rich and diverse group that needs local leaders providing inspiration. Who has the greatest potential? We examined cities with the highest COOLCITY Index values across different size categories in various regions.

Individual leaders in the regions achieve scores from 5.3 in Southern Europe up to 8.5 in Central-Eastern Europe. This is the highest score among all cities and across Europe, achieved by Piwniczna-Zdrój – a very small mountain town in Poland. Large metropolises are led by Stockholm. COOLCITY regional leaders possess high-quality resources to build nature-based resilience and become drivers of change and inspiration in their regions.

	Northern Europe	Western Europe	Central-Eastern Europe	Southern Europe
VERY LARGE > 500k people	Stockholm C 6.7	Hamburg C 6.3	Kraków C 6.0	Genoa C 5.3
LARGE 100k - 500k people	Jyväskylä C 7.4	Bergisch Gladbach Koblenz C 6.8	Rybnik C 6.8	Trento C 6.0
MEDIUM 20k - 100k people	Rudersdal Torslanda C 7.7	Plettenberg Sundern C 7.6	Konstancin-Jeziorna C 7.4	Eibar C 7.3
SMALL 10k - 20k people	Lempäälä, Nödinge-Nol C 8.0	Boppard C 8.2	Boguszów-Gorce C 7.7	Viveiro C 7.7
VERY SMALL < 10k people	li C 8.3	Bad Herrenalb C 8.1	Piwniczna-Zdrój C 8.5	Echinos C 8.4



Click on a city's name to view its profile and similar cities

Regional leaders Performance Index

How we can jointly build resilient cities?

Regional leaders in terms of the Performance Index (PI) are cities that most efficiently develop their blue-green infrastructure compared to cities with similar conditions. Although their absolute adaptive potential measured by the CCI indicator is not the highest in the regions, these towns utilise their BGI resources better than would be expected from their geographical or demographic conditions. Cities such as Basiglio (PI +1.5), Plopieni (PI +1.4), and L'île Saint-Denis (PI +1.3) perform better than similar centres, indicating effective spatial policies and appropriate use of organisational and financial resources for BGI development by their local governments.

	Northern Europe	Western Europe	Central-Eastern Europe	Southern Europe
VERY LARGE > 500k people	Stockholm +0.3	Cologne +0.3	Wrocław +0.2	Genoa +0.3
LARGE 100k - 500k people	Jönköping +0.3	Erlangen +0.6	Zielona Góra +0.7	Santa Coloma de Gramenet +0.8
MEDIUM 20k - 100k people	Hillerød +1.0	Epe +1.2	Ajka +1.2	Chaidari +1.4
SMALL 10k - 20k people	Lillerød +0.9	Ville-d'Avray +1.3	Chodov +1.1	Santa Flavia +1.4
VERY SMALL < 10k people	Trakai +0.8	Glücksburg (Baltic Sea) +1.2	Poniatowa +1.4	Basiglio +1.5




Best practices from the blue-green backyard

What actions are regional leaders taking?

Nature-based adaptation includes solutions such as developing a coherent network of green spaces in the city, planting street trees, green roofs and walls, as well as creating rain gardens and retention basins or restoring watercourses. Equally important is protecting existing blue-green infrastructure resources, including valuable wetlands. These actions reduce the urban heat island effect, improve water retention to lower flood risk, and increase ecosystem resilience and biodiversity. The range of solutions is broad, and the choice depends on local conditions and resources.

Let's look at example of actions taken by cities with the highest CCI scores in different European regions, which contribute to their strong climate resilience.



Blog case study

'ROYAL SEAPORT' DISTRICT

STOCKHOLM  Northern Europe

 Very Large City: 1,830,764 inhabitants



Stockholm Royal Seaport is a climate-resilient port district project. It is based on high standards for green spaces (Green Space Index) and stormwater management strategies. By 2024, over 14 hectares of green areas have been created here, with every resident having access to a park within 200 metres of their home. Rainwater is treated as a resource, stored in rain gardens with limited infiltration into the ground. The district also serves as an ecological corridor.

Similar cities: Göteborg, Espoo, Helsinki, Vilnius, Berlin



GREEN ROOF STRATEGY

HAMBURG  Western Europe

 Very Large City: 1,822,082 inhabitants



In 2014, Hamburg introduced a comprehensive green roof strategy for the city. The strategy includes attractive subsidy programmes for residents and businesses, awareness campaigns, planning regulations, and scientific support from HafenCity Universität Hamburg. Its goal is to cover at least 70% of suitable roofs with vegetation (100 ha). Researchers have documented the benefits: cooling the area, reducing stormwater runoff (by up to 74%), lowering flood risk, and increasing biodiversity.

Similar cities: Rīga, Warsaw, Essen, Cologne, Gdańsk

Best practices from the blue-green backyard

What actions are regional leaders taking?



BLOOMING BAD HERRENALB

BAD HERRENALB  Western Europe

 Very Small Town: 5,375 Inhabitants



This spa town located in the Schwarzwald Mitte/Nor Nature Park is implementing the “Blühendes Bad Herrenalb” (Blooming Bad Herrenalb) initiative in cooperation with the park and residents. The project aims to support biodiversity and create natural habitats for pollinating insects by transforming urban lawns into flowering meadows. It also includes educational activities promoting the creation of flower beds in gardens.

Similar towns: Recoaro Terme, Ruhla, Dobšíná, Szczyrk



CHYTEJ CO NAPADZE

RYBNIK  Central-Eastern Europe

 Large City: 125,117 Inhabitants



Rybnik, as a member of the COOLCITY project advisory team, is an example of a post-industrial city actively working towards sustainable transformation and adaptation based on the BGI. The city focuses on local rainwater retention and reducing pressure on the sewage system. Environmental education campaigns are conducted alongside a subsidy program for residents called “Chytej co napadze” to create retention basins and rain gardens.

Similar towns: Wałbrzych, Bergisch Gladbach, Borås



BIO VALUE

TRENTO  Southern Europe

 Large City: 114,302 Inhabitants



Trento incorporates the value of nature and ecosystem services into its policies. In the BioValue project, it is transforming the Fersina River from a regulated channel into an accessible socio-ecological space that supports biodiversity and city resilience while engaging residents in co-design. Plans include riverbank renaturation, introduction of riparian vegetation, linear parks along the valley, and increased floodplain areas.

Similar towns: Saarbrücken, Žilina, Moncalieri, Maribor

Best practices from the blue-green backyard

What actions are regional leaders taking?



PARCO AGRICOLO SUD MILANO

BASIGLIO  Southern Europe

 Very small town: 7,071 inhabitants



Basiglio is a municipality entirely within the Parco Agricolo Sud Milano — a vast agricultural and natural park established to protect farming, the environment, and rural landscape. The historic irrigation network forms one of the most productive agricultural structures in Europe.

Similar towns: **Blomberg, Asolo, Pont-Évêque, Hadamar**



RIVER RENATURATION

li  Northern Europe

 Very small town: 3,343 inhabitants



This is an example of a climatically active small municipality in Europe. The municipality of li carries out river renaturation projects for the Kuivajoki and Olhavanjoki rivers, and also participates in the Iijoki Old Riverbed Development Project, aimed at restoring the old Iijoki riverbed and improving fish migration conditions. Additionally, as a signatory of the Iijoki River Agreement, li is involved in activities including the restoration of drained peatlands and wetlands and the renaturation of streams throughout the Iijoki catchment area.

Similar towns: **Masku, Ādaži, Elva, Kramfors**

A place for your city's
project on the
COOLCITY blog

Submit a project





Shared challenges and adaptation opportunities

How to find similar cities and build partnerships?

The COOLCITY Index helps identify cities that are leaders in their category and have the best blue-green infrastructure resources for adaptation, as well as those facing similar challenges with comparable development conditions. This makes it not only a diagnostic tool but also a platform supporting the exchange of knowledge and best practices among local governments.

On the coolcity.eu platform, every city can find a list of similar cities—places with comparable geographical and demographic conditions and similar adaptive potential based on BGI resources. Comparisons with cities of a similar nature can be more valuable than referencing the largest metropolises, as they consider the actual capabilities and needs of the specific entity.

We encourage using the platform to establish cooperation, seek partners for experience exchange, and build a network of cities working together on climate change adaptation strategies.



Find your city on the coolcity.eu platform



Check which cities have similar conditions: geographical · demographic · adaptive



Build a network for experience exchange and act together!

COOLCITY Index in EU countries

How does adaptation potential vary across different countries?

European cities develop under diverse environmental, spatial, and social conditions, so their capacity to adapt to climate change is not uniform. Regardless of the starting point, it is crucial for local and national decision-makers to identify their own resources and manage adaptation based on measurable data.

European climate law requires member states to systematically strengthen resilience and monitor progress in implementing adaptation measures. To this end, the COOLCITY Index report and **platform** present indicators monitoring changes in urban ecosystems as outlined in European regulations [19, 20]. These indicators should be incorporated into strategic documents at both national and local levels. They include: the share of green areas in the city, tall vegetation coverage (tree canopies), percentage of impermeable surfaces, and indicators related to rainwater infiltration and urban heat.

The next section of the report features profiles of 27 European Union countries, presenting an overall picture of the environmental adaptation potential of these states, as well as the opportunities and challenges linked to local green infrastructure resources. The profiles highlight national leaders with the highest COOLCITY Index and Performance Index values, and also present indicators monitoring the state of the urban environment.


Northern Europe	Western Europe	Central Eastern Europe	Southern Europe
Denmark 49	France 60	Poland 73	Portugal 89
Sweden 51	Belgium 62	Czech Republic 75	Spain 91
Finland 53	Netherlands 64	Slovakia 77	Italy 93
Estonia 55	Luxembourg 66	Hungary 79	Greece 95
Latvia 56	Germany 67	Romania 81	Cyprus 97
Lithuania 58	Austria 69	Bulgaria 83	Malta 99
	Ireland 71	Croatia 85	
		Slovenia 87	

How to read the EU countries' cards?

Each country card includes:

- The number of analyzed urban areas, their area and population
- The adaptive potential index of the country's cities relative to the EU average
- Leaders of the **COOLCITY Index** and **Performance Index** in different city size groups
- BGI monitoring indicators for strategic documents
- Links to city profiles on the **COOLCITY Index platform** – click on a city name to view its characteristics and similar cities
- Variation in indicator values across cities grouped by population size*

HOW TO READ THE DEVIATION CHART?



Reference point



The 0 axis indicates the average indicator values in the country

Label and interpretation

The label indicates the value of the group's difference from the average of all cities in the country. The colour shows how to interpret it (positively or negatively)

Direction of deviation and strength of difference

The direction of the bar shows whether the result is above or below the average. Its length is the z-score, meaning the longer the bar, the stronger the deviation from the average



*In selected countries, variation is not shown due to too few urban areas

Denmark

EU region
Northern Europe

Number of urban areas
in COOLCITY
125

Urban area
2 141 km²

Population in urban areas
3.8 mln

COOLCITY Index

6.2

EU average: 5.5 Region average: 6.6

COUNTRY LEADERS

COOLCITY
INDEX

PERFORMANCE
INDEX

V. LARGE, LARGE
> 100k people
5 cities

Odense
6.0

Copenhagen
+0.3

MEDIUM
20 - 100 k people
39 cities

Hillerød
7.3

Hillerød
+1.0

SMALL
10 - 20 k people
31 cities


Rudersdal
7.7

Lillerød
+0.9

VERY SMALL
< 10 k people
50 cities

Fredensborg
7.6

Fredensborg
Sorø, Værløse
+0.8



COOLCITY Index




INDICATORS FOR STRATEGIC DOCUMENTS • country average

Vegetation cover	High vegetation cover	Green space area per capita	Ecological connectivity	Share of sealed surfaces	Sealed surface area per capita
72 %	17 %	608 m²/ person	78 %	28 %	228 m²/ person

Denmark

City populations and environmental conditions



Conclusions

Danish cities have a balanced **adaptive potential**.

Average **COOLCITY Index** values ranging from 6.0 to 6.3 indicate that both large and small cities have favourable resources for adaptation using BGI.

Danish cities maintain a favourable relationship between greenery and **Surface permeability**.

With vegetation cover around 70% and a permeability index above 7, they maintain a good spatial distribution for BGI.

Denmark's stable **adaptive potential** does not negate local thermal differences.

In larger cities, surface temperature is on average 1.1°C higher than the country average, increasing the importance of BGI's cooling functions.

Sweden

EU Region
Northern Europe

Number of urban areas
in COOLCITY
212

Urban area
3,984 km²

Population in urban areas
7.2 million

COOLCITY Index

6.6

EU average: 5.5 Region average: 6.6

COUNTRY LEADERS

COOLCITY
INDEX

↑
PERFORMANCE
INDEX

• VERY LARGE,
LARGE
> 100k people
10 cities

• Jönköping
7.0

• Stockholm, Jönköping
+0.3

• MEDIUM
20k - 100k people
50 cities

• Torslanda
7.7

• Torslanda
+0.7

• SMALL
10k - 20k people
63 cities


• Nödinge-Nol
8.0

• Knivsta, Skoghall, Onsala
+0.6

• VERY SMALL
> 10k people
89 cities

• Mullsjö
7.8

• Löddeköpinge
+0.7



INDICATORS FOR STRATEGIC DOCUMENTS • country average



Vegetation
cover

76 %



High vegetation
cover

30 %



Green space area per
capita

613 m²/
person



Ecological
connectivity

74 %



Share of
sealed surfaces

23 %




Sealed area
per capita

189 m²/
person

Sweden

City populations and their environmental conditions



Conclusions

Sweden has a very balanced **adaptive potential** across its cities.

Having average **COOLCITY Index** values range from 6.5 to 6.7, different city size groups have similarly favourable starting conditions for developing high-quality NBS and building climate resilience.

Very small Swedish cities stand out with higher **biodiversity**.

In this group, **biodiversity** score averages 6.6 and remains 0.2 above the country average, supporting natural values of NBS.

In larger Swedish cities, local overheating hotspots are evident.

Despite similar conditions for NBS development in Swedish cities, the largest centres show a higher share of areas with elevated temperatures (0.4 percentage points above the average).

Finland

EU Region
Northern Europe

Number of urban areas
in COOLCITY
121

Urban area
2,945 km²

Population in urban areas
3.7 million

COOLCITY Index

7.1

EU average: 5.5 Region average: 6.6

COOLCITY
INDEX

PERFORMANCE
INDEX

LEADERS IN THE COUNTRY

VERY LARGE,
LARGE
> 100k people
8 cities

Jyväskylä
7.4

Jyväskylä, Oulu
+0.2

MEDIUM
20k - 100k people
30 cities

Imatra,
Lohja
7.6

Joensuu, Kokkola
Lohja, Mikkeli
+0.3

SMALL
10k - 20k people
22 cities


Lempäälä
8.0

Valkeakoski
+0.5

VERY SMALL
< 10k people
61 cities

li
8.3

Masku
+0.7



INDICATORS FOR STRATEGIC DOCUMENTS • country average

Vegetation
cover

82 %

High vegetation
cover

35 %

Green space area per
capita

996 m²/
person

Ecological
connectivity

76 %

Share of
sealed surfaces


17 %

Sealed area
per capita

204 m²/
person

Finland

City populations and their environmental conditions



Conclusions

Finland has a balanced and stable adaptive capacity in its cities.

The average **COOLCITY Index** value is 7.1, with a standard deviation of 0.5, indicating a similar distribution of results regardless of city size category.

Waters and wetlands enhance the adaptive conditions of small Finnish cities.

Their share averages 8%, supporting retention and microclimate within the urban structure, resulting in a higher **water conditions** index (0.6 above the country average).

Higher surface temperatures are notable in Finland's large cities.

The average for this group is 1.3°C higher than the country average, indicating the presence of local heat islands in these cities and the need to apply green infrastructure to cool them down.


Estonia

EU Region
Northern Europe

Urban area
299 km²

Number of urban areas
in COOLCITY
22


Population in urban area
784k



LEADERS IN THE COUNTRY

COOLCITY INDEX **PERFORMANCE INDEX**

LARGE, MEDIUM 20 - 100k people 5 cities	Kohtla-Järve, Narva 6.4	Narva +0.5
SMALL 10 - 20k people 5 cities	Kuressaare 7.2	Kuressaare +0.4
VERY SMALL <10k people 12 cities	Elva 7.7	Elva +0.5



INDICATORS FOR STRATEGIC DOCUMENTS • country average

 Vegetation cover 77 %	 High vegetation cover 25 %	 Green area per inhabitant 534 m²/person	 Ecological connectivity 75 %	 Share of sealed surfaces 22 %	 Sealed area per capita 138 m²/person
------------------------------	-----------------------------------	--	-------------------------------------	--------------------------------------	---

Latvia

EU Region
Northern Europe

Urban area
557 km²

Number of urban areas
in COOLCITY
38

Population in urban area
1.2 million

COOLCITY Index

7.0

EU average: 5.5 Region average: 6.6

COUNTRY LEADERS

COOLCITY
INDEX

PERFORMANCE
INDEX

VERY LARGE,
MEDIUM
20 - 100k people
8 cities

Jūrmala
7.7

Jūrmala
+0.6

SMALL
10 - 20k people
9 cities


Baloži
7.9

Olaine
+0.6

VERY SMALL
> 10k people
21 cities

Ādaži
8.0

Aizkraukle
+0.6



INDICATORS FOR STRATEGIC DOCUMENTS • country average

Vegetation
cover

80 %

High vegetation
cover

29 %

Green area
per inhabitant

600 m²/
person

Ecological
connectivity

62 %

Share of
sealed surfaces


19 %

Sealed area
per capita

143 m²/
person

Latvia

City populations and their environmental conditions



Conclusions

In Latvia, the **COOLCITY Index** remains high, with moderate differences between city groups.

Average values from 6.5 to 7.2 show a stable foundation for building resilience based on the BGI index in the country.

Cities with 10k to 20k inhabitants in Latvia are notable for high **biodiversity**.

In this group, it is 7.0 and remains 0.4 above the country average, strengthening the natural potential of the BGI.

Vegetation state is lower in larger Latvian cities.

The average share of high-vigour vegetation is 76.0% and remains 6.7 percentage points below the country average, highlighting the need to strengthen the BGI in the largest cities.



Lithuania

EU Region
Northern Europe

Number of urban areas
in COOLCITY
49

Urban area
811 km²

Population in urban area
1.7 million

COOLCITY Index

6.4

EU average: 5.5 Region average: 6.6

LEADERS IN THE COUNTRY

COOLCITY
INDEX

PERFORMANCE
INDEX

VERY LARGE,
LARGE
> 100k people
3 cities

Vilnius
6.4

Vilnius
+0.2

MEDIUM
20k - 100k people
10 cities

Utena
6.7

Jonava, Utena
+0.2

SMALL
10k - 20k people
14 cities


Visaginas
7.4

Druskininkai
+0.6

VERY SMALL
> 10k people
22 cities

Trakai
7.9

Trakai
+0.8



COOLCITY Index



INDICATORS FOR STRATEGIC DOCUMENTS • country average

Vegetation
cover

73 %

High vegetation
cover

22 %

Green space area per
capita

465 m²/
person

Ecological
connectivity

60 %

Share of sealed
surfaces


26 %

Sealed area
per capita

167 m²/
person

Lithuania

City populations and their environmental conditions



Conclusions

Lithuania combines a high **COOLCITY Index** with a fairly consistent pattern of results across city groups.

Average CCI values between 6.0 and 6.5 indicate that cities have a stable adaptive capacity based on the BGI index regardless of size.

Cities with 10k to 20k inhabitants in Lithuania are notable for their good **vegetation state**.

The average share of high-vigour vegetation reaches 80.1% and remains 2.4 percentage points above the country average, enhancing the quality of green infrastructure.

Thermal conditions are less favourable in larger Lithuanian cities.

The average surface temperature is 32.1°C, which is 1.4°C higher than the country average, increasing the importance of the cooling role of BGI and indicating the need to implement NBS solutions in large cities.

France

EU Region
Western Europe

Number of urban areas
in COOLCITY
2,102

Urban area
16,353 km²


Population in urban area
37.8 million

COOLCITY Index **5.3**

EU average: 5.5 Region average: 5.8

COUNTRY LEADERS

	COOLCITY INDEX	PERFORMANCE INDEX
VERY LARGE > 500k people 4 cities	Toulouse 3.8	Lyon, Paris 0.0
LARGE 100k - 500k people 37 cities	Besançon 5.2	Nice +0.5
MEDIUM 20k - 100k people 372 cities	Verrières-le-Buisson 7.2	Champs-sur-Marne, Menton, Sceaux +1.2
SMALL 10k - 20k people 461 cities	Jouy-en-Josas 7.8	L'Île-Saint-Denis, Ville-d'Avray +1.3
VERY SMALL < 10k people 1,228 cities	Mouguerre 7.9	Émerainville, Fresnes-sur-Escaut, Gretz-Armainvilliers, Saint-Jean-de-Boiseau +1.2




INDICATORS FOR STRATEGIC DOCUMENTS • country average

 Vegetation cover 66 %	 High vegetation cover 17 %	 Green space area per capita 506 m²/person	 Ecological connectivity 65 %	 Share of sealed surfaces 33 %	 Sealed surface area per capita 219 m²/person
-------------------------------------	--	---	--	---	--

France

City populations and their environmental conditions



Conclusions

Small and very small French cities have the highest **adaptive potential** in the country.

This is the most numerous group of cities, where the **COOLCITY Index** remains close to the country average, while in medium and large cities it is lower by 0.8 and 1.4 points respectively.

In the largest French cities, **vegetation vigour** and **biodiversity** support are key.

The share of high-vigour vegetation is 11.5% lower than the country average, paired with significantly lower **biodiversity**, weakening **adaptive potential** in big cities.

Better **water conditions** in small cities are not only due to the presence of **water bodies**.

Despite a smaller share of rivers and wetlands, small cities achieve good **water condition** results, showing the importance of **Surface permeability** and high soil retention capacity when thinking about **water** in cities.



Belgium

EU Region
Western Europe

Urban area
7,491 km²

Number of urban areas
in COOLCITY
486

Population in urban area
9.7 million

COOLCITY Index

6.3

EU average: 5.5 Region average: 5.8

COUNTRY LEADERS

COOLCITY
INDEX

PERFORMANCE
INDEX

VERY LARGE,
LARGE
> 100k people
6 cities

Bruges,
Charleroi, Liège
5.3

Brussels
0.0

MEDIUM
20k - 100k people
112 cities

Chaufontaine
7.1

Schoten, Wemmel
+0.8

SMALL
10k - 20k people
138 cities


Esneux
7.8

Bonheiden,
Bornem, Dison
+0.8

VERY SMALL
< 10k people
230 cities

Viroinval
8.0

La Hulpe
+1.2



COOLCITY Index

0-1 1-2 2-3 3-4 4-5 5-6 6-7 7-8 8-9 9-10

INDICATORS FOR STRATEGIC DOCUMENTS • country average


Vegetation
cover

79 %


High vegetation
cover

17 %


Green space area per
capita

965 m²/person


Ecological
connectivity

69 %


Share of sealed
surfaces


21 %


Sealed surface
area per capita

223 m²/person

Belgium

City populations and their environmental conditions



Conclusions

Belgium maintains a consistency of urban **adaptive potential** across different cities.

The average **COOLCITY Index** value ranging from 5.9 to 6.5 for the categories indicates that urban resilience can be built on BGI across different city size classes.

Very small Belgian cities stand out for **Surface permeability**.

On average, it reaches 8.8 and remains 0.4 above the country average, creating favourable conditions for BGI development in these locations.

In larger Belgian cities, **thermal conditions** are the most significant challenge.

Average surface temperature reaches 36.1°C, 1.5°C above the country average, increasing the need to implement BGI as a cooling solution for urban spaces.

Netherlands

EU Region
Western Europe

Number of urban areas
in COOLCITY
350

Urban area
7,306 km²

Population in urban area
15.8 million


COOLCITY Index

5.9

EU average: 5.5 Region average: 5.8

COUNTRY LEADERS

	COOLCITY INDEX	PERFORMANCE INDEX
VERY LARGE > 500k people 3 cities	Amsterdam 5.2	Amsterdam +0.3
LARGE 100k - 500k people 26 cities	Ede 6.1	s-Hertogenbosch +0.4
MEDIUM 20k - 100k people 209 cities	Bergen (NH.) 7.4	Epe +1.2
SMALL 10k - 20k people 86 cities	Landsmeer 7.4	Oostzaan +1.0
VERY SMALL < 10k people 26 cities	Mook en Middelaar 7.1	Bergen (L.) +0.6



INDICATORS FOR STRATEGIC DOCUMENTS • country average

Vegetation cover

70 %

High vegetation cover

13 %

Green space area per capita

437 m²/person

Ecological connectivity

64 %

Share of sealed surfaces

29 %


Sealed surface area per capita

161 m²/person



Netherlands

City populations and their environmental conditions



Conclusions

In the Netherlands, **adaptive potential** gradually decreases as city size increases.

Average CCI values are 0.2 and 0.3 higher than the country average in very small and small towns, while in the largest cities they remain 0.5 lower, highlighting more pronounced adaptive challenges in large urban centres.

Water and wetlands are a natural foundation for adaptation, especially in larger cities.

In over one-third of Dutch cities, **water** and wetland areas cover more than 3% of the surface, supporting cities' capacity to mitigate drought and intense rainfall effects.

The largest cities feature the lowest **Surface permeability**.

In this group, it is 1.1 lower than the country average, indicating more limited space for BGI development and resulting in lower **adaptive potential**.



Luxembourg

EU Region
Western Europe

Number of urban areas
in COOLCITY
10

Urban area
92 km²

Population in urban area
246k

COOLCITY Index

6.0




EU average: 5.5 Region average: 5.8

CITIES IN THE COUNTRY

COOLCITY ↑ PERFORMANCE INDEX

LARGE 100 - 500k people 1 city	Luxembourg	CCI 5.2	-0.1
	Dudelange	5.8	-0.2
	Differdange	5.7	0.0
MEDIUM 20 - 100k people 3 cities	Esch-sur-Alzette		-0.8
	Rumelange	7.0	+0.8
	Grevenmacher	6.7	0.0
VERY SMALL <10k people 6 cities	Echternach	6.5	0.0
	Diekirch	6.3	-0.3
	Ettelbruck	6.2	+0.1
	Wiltz		-0.3



INDICATORS FOR STRATEGIC DOCUMENTS • country average

 Vegetation cover 68 %	 High vegetation cover 27 %	 Green space area per capita 400 m ² /person	 Ecological connectivity 46 %	 Share of sealed surfaces 31 %	 Sealed surface area per capita 138 m ² /person
------------------------------	-----------------------------------	---	-------------------------------------	--------------------------------------	--



Germany

EU Region
Western Europe

Number of urban areas
in COOLCITY
1,680

Urban area
28,555 km²

Population in urban area
54.5 million

COOLCITY Index

6.2


EU average: 5.5 Region average: 5.8

COUNTRY LEADERS

COOLCITY
INDEX

PERFORMANCE
INDEX

VERY LARGE > 500k people 14 cities	Hamburg 6.3	Cologne +0.3
LARGE 100k - 500k people 61 cities	Bergisch Gladbach, Koblenz 6.8	Erlangen, Offenbach am Main +0.6
MEDIUM 20k - 100k people 469 cities	Plettenberg, Sundern (Sauerland) 7.6	Senden +1.0
SMALL 10k - 20k people 454 cities	Boppard 8.2	Bad Pyrmont, Ebersberg, Mölln +1.0
VERY SMALL > 10k people 682 cities	Bad Herrenalb 8.1	Glücksburg (Baltic Sea) +1.2



INDICATORS FOR STRATEGIC DOCUMENTS • country average


Vegetation
cover

77%


High vegetation
cover

21%


Green space area per
capita

636 m²/
person


Ecological
connectivity

65%


Share of sealed
surfaces


23%


Sealed surface
area per capita

177 m²/
person

Germany

City populations and their environmental conditions



Conclusions

In Germany, city size differentiates the **adaptive potential**.

Deviation of the **COOLCITY Index** from the country average ranges from +0.2 in very small towns to -0.4 in the largest, showing varied adaptive challenges depending on the size of the city.

The greatest differences concern **biodiversity** and the coherence of the natural system.

In the largest cities, **biodiversity** is 1.2 below the country average, and ecological connectivity is 39 percentage points lower, indicating a significant challenge in creating coherent BGI links in this group.

The largest cities combine lower **Surface permeability** with high **water condition** values.

Water conditions reach their highest values in very big cities, although permeability remains 0.5 below the country average. This points to the high importance of **waters** and wetlands (above 1.6 % from the average) in building resilience in these centres.

Austria

EU Region
Western Europe

Urban area
2,305 km²

Number of urban areas
in COOLCITY
243

Population in Urban area
4.8 million

COOLCITY Index **6.0**

EU average: 5.5 Region average: 5.8

COUNTRY LEADERS

VERY LARGE,
LARGE
> 100k people
5 cities

Salzburg
6.0

Salzburg
+0.4

MEDIUM
20k - 100k people
19 cities

Kapfenberg,
Klosterneuburg
6.7

Lustenau
+0.3

SMALL
10k - 20k people
44 cities


Bad Ischl
7.0

Gänserndorf,
Groß-Enzersdorf
+0.4

VERY SMALL
< 10k people
175 cities

Breitenfurt
near Vienna
7.5

Purkersdorf
+1.0



INDICATORS FOR STRATEGIC DOCUMENTS • country average


Vegetation
cover

75 %


High vegetation
cover

19 %


Green space area per
capita

771 m²/person


Ecological
connectivity

56 %


Share of sealed
surfaces

24 %





Sealed surface
area per capita

225 m²/person



Austria

City populations and their environmental conditions

	In the country	 VERY SMALL < 10k people	 SMALL 10k – 20k people	 MEDIUM AND LARGE > 20k people
Number of COOLCITY cities	243	175	44	24
Number of city inhabitants	4.8 million	0.9 mln	0.6 mln	3.4 mln
% of country population	53 %	9.8 %	6.3 %	36.6 %
	National average Standard deviation (sd)			
 COOLCITY Index	6.0 (± 0.7)			
 Surface permeability	7.9 (± 0.9)			
Biologically active surface area (%)	75.8 (± 6.7)			
 Vegetation state	5.2 (± 1.1)			
Share of high-vigor vegetation (%)	75.0 (± 8.8)			
 Biodiversity	6.2 (± 1.0)			
Ecological connectivity (%)	56.4 (± 19.4)			
 Water conditions	4.6 (± 0.9)			
Share of water and wetlands (%)	1.6 (± 3.1)			
 Thermal conditions	6.3 (± 0.7)			

Conclusions

In Austria, the highest **adaptive potential** is concentrated in very small towns.

There the **COOLCITY Index** averages 6.1, while in other city groups the results are more similar – 5.6 CCI in small towns and 5.7 in medium and large ones.

Biologically active surfaces are an important element of the spatial structure of Austrian cities.

Their share usually ranges between 69% and 83%, indicating a broad resource of areas favourable for BGI development, though it decreases with an increase in city size.

Water bodies and wetlands are an important resource for medium and large centres.

They average 2.9% of city area, which is 1.3 % higher than the average of other cities in the country, although it does not translate to higher water condition values.

Ireland

EU Region
Western Europe

Number of urban areas
in COOLCITY
81

Urban area
1,369 km²

Population in urban area
2.8 million

COOLCITY Index **6.4**

EU average: 5.5 Region average: 5.8

LEADERS IN THE COUNTRY

COOLCITY
INDEX

PERFORMANCE
INDEX

VERY LARGE, LARGE,
MEDIUM
>100k people
20 cities

Athlone
7.0

Waterford
+0.4

SMALL
10k - 20k people
23 cities

Cavan, Longford
7.4

Donabate
+1.0


VERY SMALL
>10k people
38 cities

Westport
8.1

Ballinasloe
+0.7

COOLCITY Index

0-1 1-2 2-3 3-4 4-5 5-6 6-7 7-8 8-9 9-10



INDICATORS FOR STRATEGIC DOCUMENTS • country average

Vegetation
cover

75 %

High vegetation
cover

14 %

Green space area per
capita

589 m²/
person

Ecological
connectivity

64 %

Share of sealed
surfaces


24 %

Sealed surface
area per capita

183 m²/
person

Ireland

City populations and their environmental conditions



Conclusions

Irish cities maintain a fairly consistent **adaptive potential**.

With **COOLCITY Index** averages from 6.0 to 6.6, the BGI maintains good condition across various city classes and serves as a resource for building resilience.

Very small Irish towns are notable for high **Surface permeability**.

Averaging 8.1, it remains 0.2 above the country average, supporting favourable conditions for BGI development.

In larger Irish cities, the vegetation is less **connected**.

The average ecological connectivity of vegetation stands at 51.1%, which is 13.4% below the country average, indicating a lack of cohesion in the green network and a low level of **biodiversity** in these cities.

Poland

EU Region
Central and Eastern Europe

Urban area
9,353 km²

Number of urban areas
in COOLCITY
574

Population in urban area
20.6 million


COOLCITY Index

6.2

EU average: 5.5 Region average: 5.8

COUNTRY LEADERS

	COOLCITY INDEX	PERFORMANCE INDEX
VERY LARGE > 500k people 5 cities	Krakow 6.0	Wroclaw +0.2
LARGE 100k - 500k people 29 cities	Rybnik 6.8	Zielona Gora +0.7
MEDIUM 20k - 100k people 171 cities	Andrychow 7.4	Pulawy, Tarnobrzeg +0.9
SMALL 10k - 20k people 170 cities	Boguszow-Gorce 7.7	Olecko +0.9
VERY SMALL < 10k people 199 cities	Piwniczna-Zdroj 8.5	Poniatowa +1.4







INDICATORS FOR STRATEGIC DOCUMENTS • country average

 Vegetation cover 74 %	 High vegetation cover 23 %	 Green space area per capita 511 m ² /person	 Ecological connectivity 64 %	 Share of sealed surfaces 25 %	 Sealed surface area per capita 158 m ² /person
---	--	--	--	---	---

Poland

City populations and their environmental conditions

	In the country	● VERY SMALL < 10k people	● SMALL 10k - 20k people	● MEDIUM 20k - 100k people	● LARGE > 100k people
Number of COOLCITY cities	574	199	170	171	29
City population	20.6 million	1.3 mln	2.5 mln	6.9 mln	5.4 mln
% of national population	56 %	3.6 %	6.8 %	18.9 %	12.3 %
	National average Standard deviation (sd)				
 COOLCITY Index	6.2 (± 0.6)				
 Surface permeability	7.8 (± 0.8)				
Biologically active surface area (%)	74.9 (± 6.2)				
 Vegetation state	5.4 (± 1.1)				
Green area per capita (m²/person)	510.8 (± 259.9)				
 Biodiversity	6.4 (± 0.8)				
Ecological connectivity (%)	64.0 (± 21.2)				
 Water conditions	4.7 (± 0.8)				
 Thermal conditions	6.7 (± 0.6)				
Share of low-temperature areas (%)	3.8 (± 5.1)				

Conclusions

In Poland, city size clearly differentiates **adaptive potential**.

In very small towns, the **COOLCITY Index** averages 6.5, while in large cities around 5.9, showing a change in potential with the growth of the urban centre.

Small towns strengthen the national **biodiversity** in urban areas.

In towns with fewer than 20k inhabitants, the **biodiversity** index is 6.6, compared to 5.5 in large cities, and ecological connectivity averages 71.2% – 7.2% above the country average.

In medium and large cities, Surface permeability becomes a challenge.

It is on average 0.3 lower than the country average, indicating a smaller share of spaces conducive to **biodiversity**. In these towns, the green area per capita is also significantly lower.



Czech Republic

EU Region
Central and Eastern Europe

Number of urban areas
in COOLCITY
274

Urban area
2,779 km²

Population in Urban area
5.9 million

COOLCITY Index

6.1

EU average: 5.5 Region average: 5.8

LEADERS IN THE COUNTRY

VERY LARGE,
LARGE
> 100k people
5 cities

Ostrava
6.1

Ostrava
+0.1

MEDIUM
20k - 100k people
43 cities

Orlová
7.3

Orlová
+0.9

SMALL
10k - 20k people
63 cities

Mariánské Lázně
7.2

Chodov
+1.1

VERY SMALL
< 10k people
163 cities

Nejdek
7.9

Duchcov
+1.2



INDICATORS FOR STRATEGIC DOCUMENTS • country average

Vegetation
cover

74 %

High vegetation
cover

20 %

Green space area per
capita

508 m²/
person

Ecological
connectivity

67 %

Share of sealed
surfaces

25 %


Sealed surface
area per capita

161 m²/
person



Czech Republic

City populations and their environmental conditions



Conclusions

Czech cities have a balanced **adaptive potential**.

The **COOLCITY Index** deviation from the country average is: +0.1 in very small, -0.1 in small, and -0.2 in medium and large cities, so differences between groups are minor.

In medium and large Czech cities, vegetation fragmentation is more pronounced.

About 50% of vegetation patches remain unconnected, showing the challenge of ensuring ecological continuity in these cities.

In the largest Czech cities, the share of local cool spots is low.

Areas with lower temperatures cover only 1.2% of the cities, although overall **thermal conditions** in this group remain balanced compared to other cities in the country.



Slovakia

EU Region
Central and Eastern Europe

Number of urban areas
in COOLCITY
118

Urban area
1,169 km²

Population in Urban area
2.6 million

COOLCITY Index

5.9


EU average: 5.5 Region average: 5.8

LEADERS IN THE COUNTRY

COOLCITY
INDEX

PERFORMANCE
INDEX

LARGE, MEDIUM > 20k people 30 cities	Považská Bystrica 7.0	Humenné +0.6
SMALL 10k - 20k people 34 cities	Handlová 6.9	Stupava +0.7
VERY SMALL > 10k people 54 cities	Dobšiná 7.9	Svätý Jur +0.7



INDICATORS FOR STRATEGIC DOCUMENTS • country average

 Vegetation cover 73 %	 High vegetation cover 19 %	 Green space area per capita 440 m²/person	 Ecological connectivity 57 %	 Share of sealed surfaces 27 %	 Sealed surface area per capita 151 m²/person
------------------------------	-----------------------------------	--	-------------------------------------	--------------------------------------	---



Slovakia

City populations and their environmental conditions

	In the country	● VERY SMALL < 10k people	● SMALL 10k - 20k people	● MEDIUM AND LARGE > 20k people
Number of COOLCITY cities	118	54	34	30
Number of city inhabitants	2.6 million	0.3 mln	0.5 mln	1.8 mln
% of country population	49 %	6.4 %	9.4 %	32.9 %
National average Standard deviation (sd)				
COOLCITY Index	5.9(± 0.8)			
Surface permeability	7.6(± 0.9)			
Sealed surface area per capita (m²/person)	151.2(± 47.0)			
Vegetation state	5.0(± 1.3)			
Vegetation cover (%)	72.8(± 7.0)			
Biodiversity	6.1(± 0.9)			
Ecological connectivity (%)	57.5(± 20.4)			
Water conditions	4.5(± 0.9)			
Thermal conditions	6.2(± 0.8)			
Surface temperature (°C)	3.9(± 5.2)			

Conclusions

In Slovakia, the **COOLCITY Index** is moderately varied.

Average values ranging from 5.5 to 6.1 show that **adaptive potential** using BGI varies depending on city size.

Very small Slovak cities have high permeable surface resources.

Average Surface permeability is 7.9, remaining 0.3 points above the country average, providing favourable conditions for BGI development.

In larger Slovak cities, **thermal conditions** are less favourable.

Average surface temperature at 38.5°C is 1.2°C above the country average, increasing the importance of BGI's cooling functions.



Hungary

EU Region
Central and Eastern Europe

Number of urban areas
in COOLCITY
244

Urban area
3,259 km²

Population in urban areas
6.0 million

COOLCITY Index **5.2**

EU average: 5.5 Region average: 5.8

COUNTRY LEADERS

COOLCITY
INDEX

PERFORMANCE
INDEX

VERY LARGE,
LARGE
> 100k people
7 cities

Miskolc
5.9

Szeged
+0.5

MEDIUM
20k - 100k people
50 cities

Ózd
7.1

Ajka
+1.2

SMALL
10k - 20k people
66 cities


Törökbálint
6.9

Csongrád, Kőszeg
Oroszlány, Szigethalom
Törökbálint
+0.7

VERY SMALL
> 10k people
121 cities

Nagymaros
6.9

Alsózsolca
+1.0



INDICATORS FOR STRATEGIC DOCUMENTS • country average


Vegetation
cover

67%


High vegetation
cover

12 %


Green space area per
capita

567 m²/person


Ecological
connectivity

67 %


Share of sealed
surfaces










32%


Sealed surface
area per capita

264 m²/person

Hungary

City populations and their environmental conditions

	In the country	 VERY SMALL < 10k people	 SMALL 10k – 20k people	 MEDIUM AND LARGE > 20k people
Number of COOLCITY cities	244	121	66	57
City population	6.0 mln	0.8 mln	0.9 mln	4.3 mln
% of national population	6.3 %	8.2 %	9.5 %	45.6 %
	National average Standard deviation (sd)			
 COOLCITY Index	5.2 (± 0.6)			
 Surface permeability	6.9 (± 0.8)			
Proportion of sealed surfaces (%)	32.1 (± 6.3)			
 Vegetation state	4.2 (± 0.9)			
Vegetation cover (%)	67.3 (± 6.3)			
High vegetation cover (%)	12.4 (± 8.4)			
 Biodiversity	6.1 (± 0.8)			
Ecological connectivity (%)	67.1 (± 21.1)			
 Water conditions	3.9 (± 0.8)			
 Thermal conditions	5.2 (± 0.8)			

Conclusions

Hungarian cities have a balanced **adaptive potential**.

Average **COOLCITY Index** values remain similar across all city size classes, with the largest differences concerning individual BGI elements.

Larger Hungarian cities stand out with a high share of tall vegetation.

Their share is 3.9% higher than the country average, enhancing BGI's potential to reduce overheating and improve environmental conditions.

The greatest differences concern the continuity of the natural system.

In larger cities, ecological connectivity is 10.4% below the country average, which may limit BGI coherence.

Romania

EU Region
Central and Eastern Europe

Urban area
3,364 km²

Number of urban areas
in COOLCITY
278

Population in urban areas
9.1 million

COOLCITY Index

5.5

EU average: 5.5 Region average: 5.8

LEADERS IN THE COUNTRY

COOLCITY
INDEX

PERFORMANCE
INDEX

VERY LARGE,
LARGE
> 100k people
18 cities

Cluj-Napoca, Iași
Târgu Mureș
5.3

Bacău
+0.6

MEDIUM
20k - 100k people
64 cities

Borșa
7.3

Vaslui
+0.8

SMALL
10k - 20k people
60 cities


Breaza, Sângeorz-Băi
Vișeu De Sus
6.9

Vulcan
+0.7

VERY SMALL
> 10k people
136 cities

Anina, Tismana
7.6

Plopeni
+1.4



INDICATORS FOR STRATEGIC DOCUMENTS • country average

Vegetation
cover

71%

High vegetation
cover

17%

Green space area per
capita

585 m²/person

Ecological
connectivity

57%

Share of sealed
surfaces


28%

Sealed surface
area per capita

203 m²/person

Romania

City populations and their environmental conditions



Conclusions

In Romania, the **COOLCITY Index** clearly differentiates city size classes.

Average values ranging from 4.4 to 5.8 show that BGI potential strongly depends on the size of the city. With the smallest cities showing the highest **adaptive potential**.

Very small Romanian towns stand out with a high share of biologically active surfaces.

It averages 75.4% and remains 3.9% above the country average, creating an important spatial base for BGI development.

The largest cities have the lowest vegetation cover.

It averages 57.7% compared to the country average of 70.8%, showing a smaller resource of greenery with cooling functions. With temperatures higher by 2.5 °C on average, introducing BGI here is particularly important.



Bulgaria

EU Region
Central and Eastern Europe

Urban area
1,365 km²

Number of urban areas
in COOLCITY
121

Population in Urban area
4.2 million

COOLCITY Index

5.1

EU average: 5.5 Region average: 5.8

LEADERS IN THE COUNTRY

COOLCITY
INDEX

PERFORMANCE
INDEX

VERY LARGE,
LARGE
> 100k people
6 cities

Ruse
5.0

Burgas, Ruse
+0.3

MEDIUM
20k - 100k people
29 cities

Gabrovo
7.1

Dimitrovgrad
+1.0

SMALL
10k - 20k people
29 cities


Gotse Delchev
6.5

Gotse Delchev
+1.1

VERY SMALL
< 10k people
57 cities

Dryanovo
7.1

Kavarna, Varshets
+0.8



INDICATORS FOR STRATEGIC DOCUMENTS • country average


Vegetation
cover

64 %


High vegetation
cover

17 %


Green space area per
capita

388 m²/
person


Ecological
connectivity

68 %


Share of sealed
surfaces

36 %


Sealed surface
area per capita

202 m²/
person



Bulgaria

City populations and their environmental conditions

	In the country	● VERY SMALL < 10k people	● SMALL 10 - 20k people	● MEDIUM AND LARGE > 20k people
Number of COOLCITY cities	121	57	29	35
Number of city residents	4.2 million	0.4 mln	0.4 mln	3.4 mln
% of the country's population	66 %	5.8 %	6.3 %	53.5 %
Average in the country Standard deviation (sd)				
COOLCITY Index	5.1 (± 0.8)			
Permeability of land	6.5 (± 1.0)			
Sealed surface area per capita (m2/person)	202.0 (± 77.1)			
Vegetation state	4.3 (± 1.3)			
Green area per resident (m2/person)	387.9 (± 198.2)			
Biodiversity	6.2 (± 0.9)			
Ecological connectivity (%)	67.5 (± 21.9)			
Water conditions	3.8 (± 0.9)			
Thermal conditions	4.6 (± 1.1)			

Conclusions

Bulgaria has a very balanced **adaptive potential** in its cities.

Average **COOLCITY Index** ranges from 4.9 to 5.2 showing that BGI can be a similar resource regardless of the size of the centre.

The largest Bulgarian cities have less green space per inhabitant.

On average, there are 204.7 m² of green space area per person, indicating that creating new green areas is the biggest challenge for this group of cities.

In larger cities, the vegetation is less continuous.

The average ecological Ecological connectivity is 54.4%, which is 13.1 % below the country average, indicating a deficit of green connections in building resilience using BGI.



Croatia

EU Region
Central and Eastern Europe

Number of urban areas
in COOLCITY
100

Urban area
1,205 km²

Population in Urban area
2.1 million

COOLCITY Index

5.8

EU average: 5.5 Region average: 5.8

LEADERS IN THE COUNTRY

COOLCITY
INDEX

↑
PERFORMANCE
INDEX

• **VERY LARGE,
LARGE**
> 100k people
3 cities

• **Zagreb**
5.8

• **Zagreb**
+0.1

• **MEDIUM-SIZED**
20k - 100k people
16 cities

• **Dubrovnik**
6.5

• **Pula**
+0.9

• **SMALL**
10k - 20k people
13 cities


• **Kutina, Sveta Nedelja**
6.6

• **Kutina**
+0.7

• **VERY SMALL**
< 10k people
68 cities

• **Bakar**
7.2

• **Mali Lošinj**
+1.0



INDICATORS FOR STRATEGIC DOCUMENTS • country average


Vegetation
cover

71%


High vegetation
cover

25%


Green space area per
capita

**746 m²/
person**


Ecological
connectivity

56%


Share of sealed
surfaces

28%


Sealed surface
area per capita

**263 m²/
person**



Croatia

City populations and their environmental conditions

	In the country	● VERY SMALL < 10k people	● SMALL 10k – 20k people	● MEDIUM AND LARGE > 20k people
Number of COOLCITY cities	100	68	13	19
City population	2.1 million	0.3 mln	0.2 mln	1.6 mln
% of national population	54%	8.3%	5.1%	41%
	National average Standard deviation (sd)			
COOLCITY Index	5.8 (± 0.7)			
Surface permeability	7.4 (± 1.0)			
Share of sealed surfaces (%)	27.9 (± 8.1)			
Vegetation state	5.2 (± 1.2)			
Vegetation cover (%)	71.2 (± 8.9)			
Biodiversity	6.3 (± 0.9)			
Ecological connectivity (%)	56.4 (± 21.1)			
Water conditions	4.6 (± 0.9)			
Thermal conditions	5.6 (± 0.7)			
Surface temperature (°C)	39.5 (± 1.6)			

Conclusions

Croatia has a moderately varied **COOLCITY Index** among city groups.

With average CCI values ranging from 5.3 to 6.0, the BGI potential of cities remains moderate, although it clearly decreases in the largest centres (-0.5 from the country average).

Very small Croatian towns maintain a high **biodiversity**.

In this group, the index averages 6.4 and remains 0.2 points above the country average, supporting the natural dimension of BGI.

Thermal conditions are more dire in larger Croatian cities.

The average surface temperature reaches 40.8°C in cities with over 20k inhabitants and is 1.2°C higher than the country average, increasing the importance of BGI's cooling role.



Slovenia

EU Region
Central and Eastern Europe

Number of urban areas
in COOLCITY
42

Urban area
584 km²

Population in Urban area
922k

COOLCITY Index

6.3

EU average: 5.5 Region average: 5.8

COUNTRY LEADERS

LARGE, MEDIUM
> 20k people
9 cities

COOLCITY
INDEX

Velenje
6.7

PERFORMANCE
INDEX

Velenje
+0.9

SMALL
10k - 20k people
12 cities


Tržič
7.4

Trbovlje
+0.5

VERY SMALL
> 10k people
21 cities

Rogaška Slatina
7.5

Rogaška Slatina
+0.6



INDICATORS FOR STRATEGIC DOCUMENTS • country average


Vegetation
cover

77 %


High vegetation
cover

33 %


Green space area per
capita

639 m²/
person


Ecological
connectivity

58 %


Share of sealed
surfaces

23 %



Sealed surface
area per capita

178 m²/
person



Slovenia

City populations and their environmental conditions



Conclusions

Slovenia maintains a fairly balanced **adaptive potential** of cities.

Average values between 5.9 and 6.4 indicate uniformly good conditions for adaptation based on BGI regardless of the size of the centre.

Very small Slovenian towns maintain a high **biodiversity**.

In this group, the index averages 6.7 and remains 0.3 above the country average, enhancing the natural potential of BGI.

Small towns stand out with the most favourable **thermal conditions**.

The average surface temperature there is 38.5°C, and the share of low-temperature areas reaches 6.3%, indicating the cooling potential of BGI.

Portugal

 EU Region
Southern Europe

 Number of urban areas
in COOLCITY
267

 Urban area
2,795 km²

 Population in urban area
5.8 million

COOLCITY Index

5.1

 EU Average: 5.5  Region average: 4.9

LEADERS IN THE COUNTRY


COOLCITY
INDEX


PERFORMANCE
INDEX


VERY LARGE,
LARGE
> 100k people
5 cities

 **Vila Nova de Gaia /
Madalena / Valadares**
4.6

 **Amadora**
+0.2


AVERAGE
20 - 100k people
74 cities

 **Carvalhos / Pedroso**
6.1

 **Vialonga**
+0.8


SMALL
10 - 20k people
62 cities


 **Sandim / Olival /
Crestuma / Lever**
6.6

 **Vila Franca de Xira**
+0.7


VERY SMALL
> 10k people
126 cities

 **Aljubarrota**
6.7

 **Eixo**
+1.0



COOLCITY Index



INDICATORS FOR STRATEGIC DOCUMENTS • country average


Vegetation
cover

63 %


High vegetation
cover

17 %


Green space area per
capita

546 m²/
person


Ecological
connectivity

56 %


Share of sealed
surfaces

36 %



Sealed surface
area per capita

264 m²/
person



Portugal

City populations and their environmental conditions



Conclusions

In Portugal, there is a strong contrast in **adaptive potential** between city groups.

The **COOLCITY Index** in very small towns is 0.4 above the country average, and in larger centres 0.6 below, indicating the greatest adaptive challenges in this group.

Surface permeability is the indicator that most differentiates cities in the country.

In larger cities, it sits 0.9 below the country average. The share of sealed areas in this group is over 11% higher than in very small towns.

The cities are also differentiated by vegetation state and environmental richness.

In cities over 20k inhabitants, vegetation cover is about 7% lower than the country average, with **biodiversity** - 0.7 lower.



Spain

EU Region
Southern Europe

Urban area
10,309 million m²

Number of urban areas
in COOLCITY
1,208


Population in Urban area
36.7 million

COOLCITY Index **4.6**

EU average: 5.5 Region average: 4.9

COUNTRY LEADERS

	COOLCITY INDEX	PERFORMANCE INDEX
VERY LARGE > 500k people 6 cities	Zaragoza 3.7	València +0.3
LARGE 100k - 500k people 52 cities	Donostia 5.9	Santa Coloma de Gramenet +0.8
MEDIUM 20k - 100k people 278 cities	Eibar 7.3	Caravaca de la Cruz +1.0
SMALL 10k - 20k people 290 cities	Viveiro 7.7	Guardamar del Segura +1.1
VERY SMALL > 10k people 582 cities	Orio 8.0	Altsasu +1.2



INDICATORS FOR STRATEGIC DOCUMENTS • country average


Vegetation
cover

59 %


High vegetation
cover

13 %


Green space area per
capita

323 m²/
person


Ecological
connectivity

69 %


Share of sealed
surfaces

38 %



Sealed surface
area per capita

173 m²/
person



Spain

City populations and their environmental conditions



Conclusions

Spain has a highly varied **adaptive potential** in its cities.

The standard deviation of the **COOLCITY Index** is 1.3, showing a wide distribution of results that reflects Spain's diverse geographical conditions.

Very small towns in Spain maintain higher **biodiversity**.

In this group, it averages 6.2 and remains 0.4 above the country average. In large cities, developing diverse BGI is a challenge, visible in particularly low ecological connectivity.

Spain's largest cities have a more sealed character.

Average Surface permeability there is 4.9, 1.3 below the country average, indicating a strong need to recover space for BGI development and improve **adaptive potential**.



Italy

EU Region
Southern Europe

Urban area
19,959 km²

Number of urban areas
in COOLCITY
1,997

Population in Urban area
41.9 million

COOLCITY Index

5.1



COUNTRY LEADERS

VERY LARGE
> 500k people
6 cities

COOLCITY
INDEX
● Genoa
5.3

PERFORMANCE
INDEX
● Genoa
+0.3

LARGE
100k - 500k people
32 cities

● Trento
6.0

● Monza
+0.7

MEDIUM
20k - 100k people
395 cities

● Bagno a Ripoli
7.1

● Mentana
+1.2

SMALL
10k - 20k people
539 cities


● Veroli
7.3

● Santa Flavia
+1.4


VERY SMALL
< 10k people
1,025 cities

● Recoaro Terme
7.9

● Basiglio
+1.5



COOLCITY Index



INDICATORS FOR STRATEGIC DOCUMENTS • country average

Vegetation
cover

67 %

High vegetation
cover

14 %

Green space area per
capita

478 m²/person

Ecological
connectivity

72 %

Share of sealed
surfaces

32 %


Sealed surface
area per capita

197 m²/person



Italy

City populations and their environmental conditions



Conclusions

In Italy, the **COOLCITY Index** changes depending on the size of cities.

Average **adaptive potential** values vary by city size category – from 4.5 in the largest centres to 5.4 in the smallest with the best BGI conditions.

Very small towns in Italy have more permeable surfaces.

Average Surface permeability reaches 7.3 and remains 0.4 points above the country average, supporting the spatial foundation for BGI development.

In Italy's largest cities, vegetation is less continuous, lowering **biodiversity**.

Average ecological connectivity is 46.2%, remaining 25.6% below the country average, highlighting the challenge of building BGI networks in this city size group.



Greece

EU Region
Southern Europe

Number of urban areas
in COOLCITY
260

Urban area
3,059 km²

Population in urban area
7.8 million

COOLCITY Index **4.6**




EU average: 5.5 Region average: 4.9

LEADERS IN THE COUNTRY

COOLCITY
INDEX

↑
PERFORMANCE
INDEX

VERY LARGE, LARGE > 100k people 12 cities	Patras 4.3	Heraklion, Larissa 0.0
MEDIUM 20k - 100k people 94 cities	Arta, Corfu, Kastoria 6.0	Chaidari +1.4
SMALL 10k - 20k people 39 cities	Eleftheroupoli 6.5	Grebena +0.7
VERY SMALL > 10k people 115 cities	Echinos 8.4	Echinos +1.2



COOLCITY Index



INDICATORS FOR STRATEGIC DOCUMENTS • country average


Vegetation
cover

60 %


High vegetation
cover

13 %


Green space area per
capita

588 m²/person


Ecological
connectivity

70 %


Share of sealed
surfaces

38 %



Sealed surface
area per capita

268 m²/person



Greece

City populations and their environmental conditions



Conclusions

In Greece, the **COOLCITY Index** clearly differs between city size classes.

CCI values from 2.5 to 5.4 show that BGI potential strongly depends on the local context. In the largest centres, **adaptive potential** drops by 2 points below the country average.

Very small towns in Greece stand out for their permeable surface coverage.

Surface permeability there averages 7.3 and remains 1.2 above the country average, providing a good basis for BGI solutions.

Biodiversity is one of the main challenges in the country's largest cities.

The average index value remains 2.7 below the country average, and areas with potentially high **biodiversity** cover only 32.2% of the surface, reducing the natural potential of BGI.



Cyprus

EU Region
Southern Europe

Number of urban areas
in COOLCITY
33

Urban area
324 km²

Population in urban area
654k

COOLCITY Index

2.1

EU average: 5.5 Region average: 4.9

COUNTRY LEADERS

COOLCITY
INDEX

PERFORMANCE
INDEX

LARGE, MEDIUM
> 20k people
10 cities

Aglantzia
3.1

Aglantzia
+1.1

SMALL
10 - 20k people
8 cities


Paralimni, Ypsonas
2.6

Agios Dom
+0.5

VERY SMALL
< 10k people
15 cities

Dromolaxi
3.4

Dromolaxi
+0.6



COOLCITY Index




INDICATORS FOR STRATEGIC DOCUMENTS • country average

Vegetation cover	High vegetation cover	Green space area per capita	Ecological connectivity	Share of sealed surfaces	Sealed surface area per capita
31 %	1 %	234 m²/ person	47 %	62 %	411 m²/ person



Cyprus

City populations and their environmental conditions



Conclusions

In Cyprus, the variation in the **COOLCITY Index** divides the urban areas into three groups.

Adaptive potential is lowest in the largest centres, at only 1.6 CCI, increasing to 2.5 in very small towns. The BGI resource layout is therefore size dependent.

Very small towns in Cyprus are an important **biodiversity** reservoir.

In this group, it averages 4.3, exceeding the country average by 0.7, which strengthens the natural dimension of BGI.

In larger Cypriot cities, **Surface permeability** is lower.

It averages 2.2, which is 0.8 points below the country average, showing that space for BGI development is limited and should be increased.

Malta

EU Region
Southern Europe

Number of urban areas
in COOLCITY
6

Urban area
165 km²

Population in Urban area
478k

COOLCITY Index

3.8

EU average: 5.5 Region average: 4.9

CITIES IN THE COUNTRY

LARGE
100 - 500k people
2 cities

Saint Paul's Bay
Qormi

4.1
+0.3
+0.3

MEDIUM
20 - 100k people
4 cities

Victoria
Rabat
Valletta
Birkirkara

4.7
4.6
2.9
+0.1
0.4
-0.1
-0.2



INDICATORS FOR STRATEGIC DOCUMENTS • country average

Vegetation cover	High vegetation cover	Green space area per capita	Ecological connectivity	Share of sealed surfaces	Sealed surface area per capita
54 %	2 %	226 m ² /person	55 %	40 %	150 m ² /person

Wrocław


The First COOLCITY in Europe

Wrocław is the first city in Europe to use the full range of innovative COOLCITY tools: **COOLCITY Index**, **COOLCITY View**, and **COOLCITY Design**.



As the demonstration city for the Life project Wrocław has helped create and test solutions to meet real urban management needs from the very start.

COOLCITY provides comprehensive information about the urban environment and equips the city with practical tools to support decision-making. This enables officials to effectively build the city's resilience to climate change and improve residents' quality of life through nature-based solutions.



Wrocław in the COOLCITY Index 2026

Lower Silesian Voivodeship, Poland

BASIC INFORMATION ABOUT THE CITY



Population in urban area
663,581 people



Average total rainfall
761 mm



Urban area
195.99 km²



Average annual temperature
8.9°C




Population density
3,386 people per km²




Average total rainfall
119.53 m

IN CITY RANKINGS


	 Wrocław miasto, spółka	 Cities in Lower Silesian Voivodeship	 Very large cities in Poland	 Very large cities in EU Central-East
	Wrocław	Averages for 63 cities	Averages for 5 cities	Averages for 10 cities
 COOLCITY Index	5.8	6.3	5.8	5.8
 Surface permeability	7.5	8.0	7.4	7.5
 Vegetation state	5.4	5.7	5.3	5.0
 Biodiversity	5.2	6.6	5.4	6.3
 Water conditions	5.0	4.6	4.8	4.4
 Thermal conditions	6.0	6.6	6.1	6.0



LAND COVER



SIMILAR CITIES




They are similar in terms of:

Geographical conditions	Demographics	Adaptation challenges
 8.0 - 9.1°C		 7.3 - 8.0
 81 - 270 m a. s. l.	541,900 - 1,288,216 people	 4.9 - 5.5
 689 - 959 mm		 4.7 - 5.3
		 4.7 - 5.2
		 5.0 - 6.2

Wrocław

Potential of the city's blue-green infrastructure



Wrocław achieves a favourable surface permeability result, +0.4 higher than expected from the city's conditions. 72% of the urban area remains biologically active, preserving ample space for investments based on blue-green infrastructure.

Wrocław's space features a vegetation state index value +0.5 higher than expected based on its conditions. This shows that greenery is an important component of the city's adaptive potential, and its vigor is high – above every other very large city in Poland.

A biodiversity level of 5.2 is favourable compared to similar cities. Over 60% of urban areas have potentially high biodiversity with room for improvement in ecological connectivity. The 11% value is below average in very large Polish cities.

The city's water condition score is also higher compared to similar cities. An important natural resource is the extensive water network, which covers around 3% of its surface. These areas support fairly favourable water conditions in Wrocław. Overall, nearly 60% of the city's area has potential for rainwater retention and enhancing drought resilience.

Thermal conditions are less favourable compared to similar cities. The result is lowered by the average surface temperature of 37°C, which is 0.4 degrees higher than the average for other very large cities in the country. Areas with high temperature make up 4% of Wrocław's surface. These are areas in the city where blue-green infrastructure should be implemented to provide cooling.



Nature-based adaptation

How is Wrocław implementing NBS?

The first step to effective data-driven management is a thorough understanding of your natural resources. A comprehensive inventory of the city's environmental components allows for assessment indicating which areas have the greatest adaptation potential and which require improvement.

With the support of the COOLCITY View tool, Wrocław can make strategic decisions and set priorities and directions for action. Using spatial valuation of the city in terms of blue-green infrastructure, the city develops strategic documents such as the Urban Adaptation Plan.

Explore the COOLCITY View for Wrocław.




Surface permeability




Vegetation state



Biodiversity



Water conditions




Thermal conditions

Data-driven management

How does COOLCITY support decision-making?

The COOLCITY Design decision support system enables the city to plan implementations from a catalogue of nature-based solutions (NBS). Actions may include protection, maintenance changes, or introducing new elements of blue-green infrastructure. The system's recommendations consider both environmental benefits and residents' preferences.

COOLCITY DESIGN DECISION SUPPORT SYSTEM OPERATION SCHEME



Comparative analysis based on CCI

In the above example, the rain garden would be the winning solution

As part of the Life COOLCITY project, Wrocław will conduct pilot interventions aimed at improving local nature. By monitoring adaptive indicator changes, the city will be able to demonstrate the effectiveness of implemented NBS solutions both during the pilot and across the entire urban area.

“

The Life COOLCITY project provides Wrocław primarily with knowledge and professional tools to better manage greenery and urban space.

Thanks to modern technologies, we can precisely analyse the environmental condition and make targeted decisions that genuinely improve residents' quality of life. This tool will allow us not only to monitor changes but also to effectively counteract the negative effects of urbanisation and climate change by placing greenery where it will have the greatest impact in minimising the urban heat island effect and enhancing quality of life.



Katarzyna Szymczak-Pomianowska

Director of the Strategy and Sustainable Development Department
Wrocław City Hall

“

COOLCITY Design includes a decision support system that allows us to model various scenarios for implementing blue-green infrastructure in the urban space. This helps us identify priority areas for action and determine the specific measures to implement to maximise environmental and social benefits.



Małgorzata Brykarz

Director of the Climate and Energy Department
Wrocław City Hall

Methodology

A city in COOLCITY

There is no universal definition of a city in the European Union. In COOLCITY we define cities as an urban areas managed by a single administrative authority.

To define the city area, we use two sets of input data:

- administrative division LAU (Local Administrative Units)
- degree of urbanisation DEGURBA, which considers population size, population density, and building density
- a point layer of cities supplemented with population data obtained from Eurostat

Based on this data, we identify urban areas that must meet specific demographic and area criteria:

- a population of at least 5,000 inhabitants according to local statistical offices
- an urban area size of no less than 2 km². Cities located in agglomerations are exempt from the area size criterion.

Ultimately, we found 11,041 urban areas in the European Union that meet these criteria.

INITIAL DATASET (LAU AND DEGURBA)



SELECTION OF LAU AREAS WITH URBAN AREA > 2 KM²



SELECTION OF LAU AREAS MEETING POPULATION THRESHOLD > 5,000 INHABITANTS



CLIPPING URBAN AREAS TO LAU BOUNDARIES



COOLCITY URBAN AREAS



- Local administrative units (LAU)
- + Urban areas DEGURBA
- Selected areas
- Discarded areas
- COOLCITY urban areas

Methodology

How is the COOLCITY Index (CCI) created?


The COOLCITY Index platform and report are published annually and include data current for the previous calendar year.

Initially, we developed a set of thematic maps: land cover, vegetation vigour, surface temperature, and surface radiative properties. These provide a consistent source of spatial data for cities across the European Union and are based on satellite data from Sentinel-2, Landsat-8, Landsat-9, and Google Dynamic World. From these maps, we calculated 15 environmental indicators for each urban area (one indicator equals one value per city).

Next, we grouped and combined the indicators using a weighted average into 5 adaptive challenge indicators. These represent the state of a city's blue-green infrastructure in terms of key urban natural aspects: surface permeability, vegetation state, biodiversity, thermal conditions, and water conditions.

Finally, we calculated the COOLCITY Index (CCI) value as the average of the 5 adaptive challenge indicators. The CCI is a synthetic indicator of a city's adaptive potential, used to assess the state of blue-green infrastructure within urban areas.

The result is a diagnosis of 11,041 European cities regarding their environmental adaptive potential. Lower CCI values indicate lower potential and signal greater adaptive challenges, while higher values indicate greater potential to strengthen city resilience based on blue-green infrastructure.



Indicators

Complete list of maps and environmental indicators

Adaptive challenge indicators
forming the COOLCITY Index

Environmental indicators

Description

Thematic maps

Surface permeability

Share of sealed surfaces [%]

Percentage share of sealed surfaces in
the city area

Land cover map

Biologically active area [%]

Percentage share of biologically active area in
the city area

Vegetation state

Vegetation cover [%]

Percentage share of vegetation
in the city's land area

Land cover map

High vegetation cover [%]

Percentage share of high vegetation
in the city's land area

Share of high-vigour vegetation [%]

Percentage share of high-vigour vegetation
(excluding arable land)

Vegetation vigour map
Land cover map

Biodiversity

Ecological connectivity [%]

Degree of connections between vegetation patches in
the city. Describes continuity and accessibility of green
areas in the city

Land cover map

Vegetation diversity
[synthetic measure]

Degree of diversity of vegetation forms calculated
based on area and share of individual vegetation
classes. Indicator ranges from 0 to 1

Number of vegetation forms
[synthetic measure]

Average number of vegetation classes in the
reference unit. Indicator ranges from 0 to 6

Areas of high biodiversity [%]

Percentage share of areas with potentially high
biodiversity in the city area

Adaptive challenge indicators forming the COOLCITY Index

Environmental indicators

Description

Thematic maps

Water conditions	Surface retention capacity [%]	Potential water retention capacity of the area, expressed as a percentage and estimated based on land cover classes.	Land cover map
	Share of wetland ecosystems	Percentage share of water and wetland ecosystems in the city's area	
Thermal conditions	Surface thermal properties [synthetic measure]	Indicator defining the radiative properties of the land surface in relation to solar radiation. Values range from -1 to 0. Higher values indicate favourable substrate properties (high share of lush vegetation and low albedo), while lower values indicate unfavourable substrate properties (lack of vegetation or its poor condition and high albedo)	Map of thermal properties of the substrate
	Surface temperature [°C]	City surface temperature expressed in °C	
	Share of high-temperature areas	Percentage share of areas with high temperature in the city's area	Surface temperature map
	Share of low-temperature areas	Percentage share of areas with low temperature in the city's area	

Data sources

Beck, H. E., McVicar, T. R., Vergopolan, N., Berg, A., Lutsko, N. J., Dufour, A., Zeng, Z., Jiang, X., van Dijk, A. I. J. M., & Miralles, D. G. (2023). High-resolution (1 km) Köppen–Geiger maps for 1901–2099 based on constrained CMIP6 projections. *Scientific Data*, 10, 724. <https://doi.org/10.1038/s41597-023-02549-6>

Brown, C. F., Brumby, S. P., Guzder-Williams, B., et al. (2022). Dynamic World: Near real-time global 10 m land use land cover mapping. *Scientific Data*, 9, 251. <https://doi.org/10.1038/s41597-022-01307-4>

Copernicus Data Space Ecosystem (2021). Copernicus DEM GLO-30 — Global Digital Elevation Model. <https://doi.org/10.5270/ESA-c5d3d65>

Drusch, M., Del Bello, U., Carlier, S., Colin, O., Fernandez, V., Gascon, F., ... Bargellini, P. (2012). Sentinel-2: ESA's optical high-resolution mission for GMES operational services. *Remote Sensing of Environment*, 120, 25–36. <https://doi.org/10.1016/j.rse.2011.11.026>

Earth Resources Observation and Science (EROS) Center. (2020). Landsat 8–9 Operational Land Imager / Thermal Infrared Sensor Level-2, Collection 2 [Data set]. U.S. Geological Survey. <https://doi.org/10.5066/P9OGBGM6>

European Space Agency (ESA). (2015–present). Sentinel-2 MSI: MultiSpectral Instrument, Level-2A [Data set]. Google Earth Engine. https://developers.google.com/earth-engine/datasets/catalog/COPERNICUS_S2_SR_HARMONIZED

Eurostat. (2021). NUTS 2021 administrative boundaries. GISCO, European Commission, <https://ec.europa.eu/eurostat/web/gisco/geodata/statistical-units/territorial-units-statistics>

Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., & Moore, R. (2017). Google Earth Engine: Planetary-scale geospatial analysis for everyone. *Remote Sensing of Environment*, 202, 18–27. <https://doi.org/10.1016/j.rse.2017.06.031>

Gutierrez, J. M., Jones, R. G., Narisma, G. T., et al. (2021). Atlas. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press. <https://doi.org/10.1017/9781009157896.021>

Hersbach, H., et al. (2018). ERA5 hourly data on single levels from 1940 to present [Data set]. Copernicus Climate Change Service, Climate Data Store. <https://doi.org/10.24381/cds.adbb2d47>

Huffman, G. J., Stocker, E. F., Bolvin, D. T., Nelkin, E. J., & Tan, J. (2023). GPM IMERG Final Precipitation L3 1 day 0.1 degree × 0.1 degree V07 [Data set]. Goddard Earth Sciences Data and Information Services Center. <https://doi.org/10.5067/GPM/IMERGDF/DAY/07>

Le Gal, M., Fernandez-Montblanc, T., Duo, E., Montes Perez, J., Cabrita, P., Souto Ceccon, P., Gastal, V., Ciavola, P., & Armaroli, C. (2023). A new European coastal flood database for low–medium intensity events. *Natural Hazards and Earth System Sciences*, 23, 3585–3602. <https://doi.org/10.5194/nhess-23-3585-2023>

Lehner, B., & Grill, G. (2013). Global river hydrography and network routing: Baseline data and new approaches to study the world's large river systems. *Hydrological Processes*, 27(15), 2171–2186. <https://doi.org/10.1002/hyp.9740>

Messenger, M. L., Lehner, B., Grill, G., Nedeva, I., & Schmitt, O. (2016). Estimating the volume and age of water stored in global lakes using a geo-statistical approach. *Nature Communications*, 7, 13603. <https://doi.org/10.1038/ncomms13603>

Pasquarella, V. J., Brown, C. F., Czerwinski, W., & Rucklidge, W. J. (2023). Comprehensive quality assessment of optical satellite imagery using weakly supervised video learning. In *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition Workshops* (pp. 2125–2135). <https://doi.org/10.1109/CVPRW59228.2023.00206>

Bibliography

[1] Nieuwenhuijsen, M. J. (2024). Climate crisis, cities, and health. *The Lancet*, 404(10463), 1693–1700. [https://doi.org/10.1016/S0140-6736\(24\)01934-2](https://doi.org/10.1016/S0140-6736(24)01934-2)

[2] Iungman, T., Khomenko, S., Barboza, E. P., Cirach, M., Gonçalves, K., Petrone, P., Erbertseder, T., Taubenböck, H., Chakraborty, T., & Nieuwenhuijsen, M. (2024). The impact of urban configuration types on urban heat islands, air pollution, CO2 emissions, and mortality in Europe: A data science approach. *The Lancet Planetary Health*, 8(7), e489–e505. [https://doi.org/10.1016/S2542-5196\(24\)00120-7](https://doi.org/10.1016/S2542-5196(24)00120-7)

[3] Perrelet, K., Moretti, M., Dietzel, A., Altermatt, F., & Cook, L. M. (2024). Engineering blue-green infrastructure for and with biodiversity in cities. *npj Urban Sustainability*, 4(1), Article 27. <https://doi.org/10.1038/s42949-024-00163-y>

[4] Li, Y., Svenning, J.-C., Zhou, W., Zhu, K., Abrams, J. F., Lenton, T. M., Ripple, W. J., Yu, Z., Teng, S. N., Dunn, R. R., & Xu, C. (2024). Green spaces provide substantial but unequal urban cooling globally. *Nature Communications*, 15, Article 7108. <https://doi.org/10.1038/s41467-024-51355-0>

[5] Kowarik, I., Fischer, L. K., Haase, D., Kabisch, N., Kleinschroth, F., Konijnendijk, C., Straka, T. M., & von Haaren, C. (2025). Promoting urban biodiversity for the benefit of people and nature. *Nature Reviews Biodiversity*, 1, 214–232. <https://doi.org/10.1038/s44358-025-00035-y>

[6] Rezaei, N., & Millard-Ball, A. (2023). Urban form and its impacts on air pollution and access to green space: A global analysis of 462 cities. *PLOS ONE*, 18(1), e0278265. <https://doi.org/10.1371/journal.pone.0278265>

[7] Santamouris, M., & Osmond, P. (2020). Increasing green infrastructure in cities: Impact on ambient temperature, air quality and heat-related mortality and morbidity. *Buildings*, 10(12), 233. <https://doi.org/10.3390/buildings10120233>

[8] Staccione, A., Candiago, S., & Mysiak, J. (2022). Mapping a green infrastructure network: A framework for spatial connectivity applied in Northern Italy. *Environmental Science & Policy*, 131, 57–67. <https://doi.org/10.1016/j.envsci.2022.01.017>

[9] Widera, B. (2023). Nature-based solutions in the context of climate-resilient buildings and cities. In V. Ramanathan & J. von Braun (Eds.), *Resilience of people and ecosystems under climate stress* (Scripta Varia 152). Pontifical Academy of Sciences. <https://www.pas.va/en/publications/scripta-varia/sv152pas/widera.html>

[10] Eurostat. (2023). Urban-rural Europe – introduction. *Statistics Explained*. https://ec.europa.eu/eurostat/statistics-explained/index.php?title=Urban-rural_Europe_-_introduction

[11] European Environment Agency. (2024). Urban. Climate-ADAPT. <https://climate-adapt.eea.europa.eu/en/eu-adaptation-policy/sector-policies/urban>

[12] World Health Organization Regional Office for Europe. (2017). Urban Green Space and Health: Intervention Impacts and Effectiveness. World Health Organization Regional Office for Europe. <https://cdn.who.int/media/docs/librariesprovider2/euro-health-topics/environment/urban-green-space-intervention.pdf>

[13] World Meteorological Organization. (2026). WMO confirms 2025 was one of warmest years on record. <https://wmo.int/news/media-centre/wmo-confirms-2025-was-one-of-warmest-years-record>

[14] European Commission. (2025). EU Climate Action Progress Report 2025. Chapter 6: Climate resilience and adaptation. Directorate-General for Climate Action. https://climate.ec.europa.eu/eu-action/climate-strategies-targets/progress-climate-action/eu-climate-action-progress-report-2025/chapter-6-climate-resilience-and-adaptation_en

[15] World Health Organization Regional Office for Europe. (2025). Climate change and health. <https://www.who.int/europe/news-room/fact-sheets/item/climate-change>

[16] European Environment Agency. (2024). European Climate Risk Assessment (EEA Report No. 01/2024). Publications Office of the European Union. <https://www.eea.europa.eu/en/analysis/publications/european-climate-risk-assessment>

[17] European Commission. (2021). EU Adaptation Strategy. https://climate.ec.europa.eu/eu-action/adaptation-and-resilience-climate-change/eu-adaptation-strategy_en

[18] European Environment Agency. (n.d.). Nature-based solutions. Climate-ADAPT. <https://climate-adapt.eea.europa.eu/en/eu-adaptation-policy/key-eu-actions/NbS>

[19] European Commission. (n.d.). Nature Restoration Regulation. https://environment.ec.europa.eu/topics/nature-and-biodiversity/nature-restoration-regulation_en

[20] European Commission. (2020). Biodiversity Strategy for 2030. https://environment.ec.europa.eu/strategy/biodiversity-strategy-2030_en

Image sources

P • 43:

Hjorthagsparken fot. www.norradjurgardsstaden2030.se
<https://www.norradjurgardsstaden2030.se/articles/green-space-index-and-stormwater-strategy>

Flakturm IV (Zielony Bunkier) fot. Tulipasyvestris, Wikimedia Commons, CC BY 4.0
https://commons.wikimedia.org/wiki/File:2024_Hamburger_Flakt%C3%BCrme_in_St._Pauli_02.jpg

P • 44:

Bad Herrenalb fot. Rick Eichner, my.toubiz, CC by SA

Fot. zbiornik na deszczówkę, www.rybnik.eu
<https://www.rybnik.eu/dla-mieszkancow/gospodarka-wodno-sciekowa/dotacje-do-retencji>

Rzeka Fersina w Trento fot. biovalue-horizon.eu
<https://biovalue-horizon.eu/>

P • 45:

Parco Agricolo Sud Milano fot. Yoruno, Wikimedia Commons, CC BY-SA 3.0.
<https://commons.wikimedia.org/wiki/File:ParcoAgricoloSudMilano.jpg>

Rzeka li fot. Ninara, flickr.com, CC BY 2.0
<https://www.flickr.com/photos/37583176@N00/2757072651/>